

HEALTH EMPIRICS

Journal of the Indian Health Economics and Policy Association
Volume 1, Issue 1, March 2026

IHEPA

Indian Health Economics and Policy Association

ISSN- 

Health Empirics

Volume 1 | Issue 1 | March 2026

ISSN:

PUBLISHER

Health Empirics is published by the Indian Health Economics and Policy Association (IHEPA).
Registered Office: Centre for Multi-Disciplinary Development Research (C.M.D.R)
Dr. B.R. Ambedkar Nagar, Lakamanahalli, Dharwad, Karnataka – 580 004, India.
Secretariat & Editorial Office: Department of Economics, Tripura University, Suryamaninagar,
West Tripura, Tripura, India, Pin- 799022.

CORRESPONDENCE

All correspondence regarding contributions, subscriptions, and association membership should be addressed to The Secretary, IHEPA at the Secretariat address above.

Email: secretary.ihepa@gmail.com | editor@healthempirics.org

COPYRIGHT & LICENSING

Copyright © 2026 The Author(s).

© All articles in Health Empirics are published under the Creative Commons Attribution 4.0 International License (CC BY 4.0). The authors retain the copyright and full publishing rights without restrictions, granting Health Empirics the right of first publication. This license permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

DISCLAIMER & ETHICS

Disclaimer: The opinions expressed in Health Empirics are those of the authors and do not necessarily reflect the views of the Indian Health Economics and Policy Association (IHEPA), the Editorial Board, or the Publisher. While the advice and information in this journal are believed to be true and accurate at the date of publication, neither the authors, the editors, nor the publisher can accept any legal responsibility for any errors or omissions that may be made.

Ethical Standard: Health Empirics adheres to the guidelines of the Committee on Publication Ethics (COPE) and the International Committee of Medical Journal Editors (ICMJE). All research involving human participants must have obtained appropriate ethical approval.

STATEMENT OF OWNERSHIP

(Form IV — See Rule 8)

Place of Publication: Department of Economics, Tripura University (A Central University), Suryamaninagar, Agartlala, Tripura – 799 022, India

Periodicity of Publication: Quarterly

Publisher's Name: Secretary, Indian Health Economics and Policy Association (IHEPA)

Editor's Name: Udaya Shankar Mishra, International Institute for Population Sciences (IIPS), Mumbai, India

Owner: Indian Health Economics and Policy Association (IHEPA)

I, Salim Shah, hereby declare that the particulars given above are true to the best of my knowledge and belief.

Sd/-
Salim Shah

Health Empirics

Journal of the Indian Health Economics and Policy Association

Volume 1, Issue 1, March 2026

Editor-in-Chief

Udaya Shankar Mishra

Editorial Board:

Amit Kumar Bardhan
Debadityuti Das
Upasak Das
Soma Dey
Saswata Ghosh
S K Godwin
William Joe
Sunil Rajpal
Mala Ramanathan
M Sivakami

Associate Editors:

Kajori Banerjee
Debasis Barik
Parma Chakravartti
Gautam Kumar Das
T R Dilip
Reshmi Sengupta
Vandana Khaitan
Abhishek Kumar
Rudra Narayan Mishra
Subrata Mukherjee
Vaibhav Puri
Sukhdeep Singh
Sunu C Thomas

Managing Editor

Salim Shah

Editorial Assistants

Komal Ahluwalia
Anirudhan P Edathil
Tulika Rohilla

Designers

Pannita Jain
Dhananjay Upadhyay

Web Developer

Gudiya Gupta

Information for Authors

Health Empirics invites submissions of Original Research, Methodological Papers, Policy Analyses, and Systematic Reviews. We particularly encourage submissions from doctoral researchers and practitioners working at the intersection of research and policy.

Submission Quick Guide:

- Manuscript Length: Recommended length is 4,000–8,000 words, including references and appendices.
- File Format: Microsoft Word (.doc, .docx) only.
- Formatting:
 - ◊ Use APA 7th Edition style for citations and references.
 - ◊ Font: Times New Roman, 12-point, 1.5 line spacing.
 - ◊ Margins: 2.5 cm (1 inch) on all sides.
 - ◊ Abstract: Structured (Background, Methods, Results, Conclusion); 200–250 words.
- Review Process: All submissions undergo an initial desk review (1 week). Manuscripts passing screening advance to double-blind peer review (approx. 6 weeks).
- Copyright & Open Access: Authors retain the copyright to their work. Articles are published under the Creative Commons Attribution 4.0 International License (CC BY 4.0), allowing for broad dissemination provided the original work is properly cited.

Fees:

Health Empirics does not charge any submission or publication fees.

How to Submit: Please email your complete manuscript and cover letter to:
editor@healthempirics.org



Scan for full Author Guidelines
and Ethics Policy.

Table of Contents

Editorial Note <i>Udaya Shankar Mishra</i>	01
The Health Factor in Multidimensional Poverty: Trends and Inequalities in India, 2005–2021 <i>Komal Ahluwalia, Anirudhan P Edathil, Saroj Kumar and William Joe</i>	03
Prevalence and Correlates of Overweight and Obesity in Adults and Older Adults in India; Population–Level Estimates Based on Nationally Representative Surveys (2015–21) <i>Debayanti Bhowmick, Abhishek Kumar and Ajay Kumar Verma</i>	30
Out-of-Pocket Health Expenditure: A Repeated Cross-Sectional Analysis of National Sample Survey Data, 2004–11 to 2022 <i>Abhishek Kumar</i>	47
Livelihood and Status of Tobacco Processing Workers: Insights from Selected States in India <i>Nayanatara S.Nayak, Rudra N. Mishra, Karabi Mujumdar, Tara Nair and N.L.Narasimha Reddy</i>	64
Productivity and Technical Change in the Indian Pharmaceutical Sector: A Comparison of Foreign and Domestic Firms <i>Tulika Rohilla and Boppana Nagarjuna</i>	82
Burden of Distress Financing for Hospitalization in India: Prevalence and Patterns from Household Health Care Consumption Survey, 2017–18 <i>Sunil Rajpal, Sneha Gupta and Shreya Ronanki</i>	98
In Memory: Dr. Mayanka Ambade (1991–2025) <i>Sunil Rajpal</i>	116

Editorial Note

Udaya Shankar Mishra*

Editor-in-Chief, Health Empirics

It is with distinct privilege that we introduce the inaugural issue of Health Empirics. This publication is realized through an overwhelming response from the research community, characterized notably by the robust submissions of emerging scholars dedicated to the rigorous inquiry on health-related topics.

The nomenclature of this journal, Health Empirics, is deliberate. In the contemporary landscape of public policy, the demand for meaningful, actionable research has shifted decisively toward the empirical. While theoretical frameworks remain foundational, the complexity of modern health challenges necessitates evidence-based governance. We posit that the most effective bridge between observation and intervention is constructed upon rigorous data analysis. Thus, this journal is established to prioritize empirical scrutiny as the bedrock of policy formulation.

Distinguished from publications that operate within the siloes of clinical science, population health, or pure economics, Health Empirics adopts a multidisciplinary mandate. We recognize that health is multifaceted, existing simultaneously as a biological reality, an economic asset, and a social indicator. By fostering the interdisciplinary convergence, we aim to elucidate complex policy connections and provide contributions that are not only academically sound but pragmatically transformative.

While the empirical evidence in this inaugural issue is primarily drawn from the Indian context, a nation navigating a profound epidemiological and demographic transition, the scope of Health Empirics extends well beyond these borders. We view India as a critical case study with significant extrapolative value for developing economies; however, our mandate is inclusive. We actively invite research contributions from the broader Global South and emerging markets to foster a comparative and comprehensive discourse on health and development.

In the current discourse on human well-being, health occupies the center stage, particularly as we strive to sustain the accomplishments of increased longevity and maximize the dividends of development. The opening analysis by Ahluwalia et al. offers a granular decomposition of India's multidimensional poverty. While the study confirms a commendable secular decline in headcount ratios, it concomitantly

* International Institute for Population Sciences (IIPS), Mumbai, India
Email: umishra@iipsindia.ac.in

exposes deep-seated persistence in nutritional deprivation, identified here as the primary structural brake on further progress. However, the escape from poverty remains precarious. In a complementary analysis of household expenditure by Rajpal et al., this issue presents compelling evidence on the fragility of financial protection. With nearly half of inpatient cases necessitating ‘distress financing’, forced borrowing, or asset liquidation, the research highlights that without universal insurance coverage, the medical poverty trap remains a potent threat to economic stability. Extending this discussion on financial vulnerability, Kumar analyzes trends in out-of-pocket and catastrophic health expenditure in India. While the share of medical spending has marginally declined, the persistence of catastrophic expenditure, particularly among SC, OBC, casual labor households, and higher consumption groups, underscores enduring gaps in financial protection despite policy efforts.

Parallel to these economic challenges, the epidemiological landscape is undergoing a profound transition. We present a study that maps the contours of this shift. At the national level, Bhowmick et al. provide new estimates of the obesity epidemic, revealing a stark gendered vulnerability and a strong correlation with urbanization, signaling that ‘diseases of affluence’ are becoming entrenched. This contribution illuminates a complex reality where the vulnerabilities of aging are increasingly intersecting with the rising tide of non-communicable diseases. Finally, we turn to the industrial and labor dynamics examined by Rohilla and Nagarjuna that undergird the health sector. The efficiency of the pharmaceutical industry is examined through Data Envelopment Analysis, where findings attribute superior technical progress in foreign firms to direct investment, linking macroeconomic policy directly to sectoral productivity. Yet, this industrial lens must not obscure the human cost of production. Another significant exploration of this issue is Nayak et al.’s assessment of the informal workforce in tobacco processing, revealing an abysmal landscape of occupational hazards. This serves as a stark reminder of the often-invisible health costs embedded within the supply chain.

This collection offers a nuanced, evidence-based outlook on the varying domains of health-related vulnerabilities. It stands as an exemplary assembly of works, indicating the potential for the scientific treatment of data to aid intervention. The evidence generated on this platform aims not merely to inform the discourse but to reform the course of action toward realizing our collective targets for a healthier society.

Finally, the realization of this inaugural issue would not have been possible without a collective effort. I extend my sincere gratitude to the Indian Health Economics and Policy Association, Editorial Board of Health Empirics, the Associate and Managing Editors, and the journal support staff for their unwavering dedication to bringing this vision to fruition.

The Health Factor in Multidimensional Poverty: Trends and Inequalities in India, 2005–2021

Komal Ahluwalia[#], Anirudhan P Edathil^{*}, Saroj Kumar^{*}, William Joe^{*}

India's commitment to Sustainable Development Goal 1—to end poverty in all its forms—necessitates a multidimensional approach beyond conventional income measures. This study investigates the evolution and determinants of multidimensional poverty in India, analyzing its incidence, intensity, and structural drivers across demographic and social groups using the Alkire–Foster framework. Using data from three National Family Health Survey (NFHS) rounds (2005–06, 2015–16, and 2019–21), the study computes the Multidimensional Poverty Index (MPI) across health, education, and living standard dimensions. Advanced decomposition techniques such as Kitagawa and Blinder–Oaxaca were employed to quantify the relative contributions of headcount ratios, poverty intensity, and socioeconomic characteristics across caste, wealth quintiles, religion, and regions. India's MPI declined markedly from 0.282 in 2005–06 to 0.066 in 2019–21, driven by reductions in the headcount ratio (52.3% to 14.7%) and moderate improvement in intensity (53.9% to 44.5%). The Scheduled Tribes, children under 14 years, rural households, and females remained the most deprived groups. Decomposition analyses indicated that 85–89% of poverty reduction stemmed from falling headcount ratios, with improvements in endowments especially education, maternal health, and access to assets playing a key role. The nutrition indicator emerged as the single largest contributor to poverty, accounting for over one-third of total deprivations. Despite remarkable progress in reducing multidimensional poverty, substantial intergroup and regional disparities persist. Strengthening nutrition, education, and social inclusion policies is essential for accelerating equitable poverty reduction and achieving sustainable human development in India.

[#]Corresponding Author: Komal Ahluwalia, Institute of Economic Growth, Delhi, India
Email: komalahluwalia2261@gmail.com

^{*}Institute of Economic Growth, Delhi, India

The United Nations' Sustainable Development Goal¹ established a mandate to end poverty in all its forms everywhere by 2030 (United Nations, 2024). Poverty is widely acknowledged as a multifaceted phenomenon that includes multiple deprivations, including but not limited to health, education, and other often overlooked dimensions. Consequently, an assessment of poverty necessitates a multidimensional approach. This multidimensional perspective on resource scarcity has gradually emerged as a critique of mainstream economic development paradigms, historically prioritizing growth in per capita Gross National Product (GNP) (Roy et al., 2019). Nevertheless, poverty analysis in many countries continues to be constrained by a unidimensional framework. These analyses rely solely on equivalent consumption as a proxy for poverty, thereby neglecting to capture the full extent of deprivations experienced by populations.

In response to these methodological limitations, scholars have proposed various refinements to poverty measurement frameworks. From the Human Poverty Index (HPI) and Human Development Index (HDI), which first introduced multidimensional perspectives in the late 1990s (Sen & Anand, 1997; UNDP, 1997) to the Multidimensional Poverty Index (MPI) developed by Alkire and Foster (Alkire & Foster, 2011; Alkire & Santos, 2011). The MPI represents a pivotal advancement, enabling direct measurement of cumulative deprivations at the household level through a dual-cutoff counting methodology for poverty classification. Unlike its predecessors, the MPI's theoretical architecture covers three dimensions operationalized through ten distinct indicators, facilitating granular analysis of both the nature and intensity of household-level deprivations. India has adopted a domestically adapted MPI developed by the NITI Aayog, adhering to the Alkire and Foster (AF) methodology employed in the global MPI. While incorporating all ten global MPI deprivation indicators, the national index expands its scope to include maternal health and bank accounts, aligning with India's development priorities (Alkire & Santos, 2011).

The Multidimensional Poverty Index offers valuable insights into the multifaceted nature of poverty beyond simple headcount measures. Existing research has explored multidimensional poverty patterns in India (Pradhan et al., 2022; Das et al., 2022), consistently demonstrating significant correlations between poverty indices and deprivation patterns among marginalized groups, particularly Scheduled Castes and Scheduled Tribes (Pradhan et al., 2022; Bagli & Tewari, 2019; Kaibarta et al., 2022), Muslim communities (Das et al., 2022) and rural inhabitants (Das et al., 2023). While existing literature has explored multidimensional poverty patterns in India, few studies have provided a long-term analysis spanning 15 years of demographic shifts. This study contributes to the literature by filling a critical methodological gap: the application of the Blinder–Oaxaca decomposition to the Multidimensional Poverty Index (MPI). Unlike previous studies that largely rely on descriptive trends or standard regression analyses, this paper systematically

disentangles the ‘endowment effects’ (changes in population characteristics) from ‘coefficient effects’ (returns to those characteristics). This offers a perspective on whether the reduction in poverty among marginalized groups (SCs/STs) is due to improved attributes or structural reductions in the disadvantages they face. This comprehensive approach not only highlights intersectional disparities but also aims to identify targeted intervention opportunities, ultimately aiming to develop more precise and effective strategies for poverty alleviation.

Methodology

The global multidimensional poverty index has undergone important methodological evolutions since its inception in 2010. Currently, the dual-cutoff counting methodology developed by (Alkire and Foster., 2011) is being utilised in the calculation of global multidimensional poverty index. The Alkire–Foster framework for measuring multidimensional poverty is grounded in Amartya Sen’s concept of capability deprivation and Atkinson’s counting methods. This methodology employs a two-step approach: identification and aggregation. The dual cut-off method involves setting thresholds for both poverty and deprivation. An individual is considered poor if their deprivation score reaches or surpasses the poverty cutoff, ‘k’. A universal cutoff of one-third ($k=0.333$) has been established. The Alkire–Foster (AF) methodology is particularly advantageous for this analysis vis-à-vis other poverty measures because of its property of subgroup decomposability. Unlike standard headcount ratios or regression residuals, the AF method allows the Multidimensional Poverty Index (MPI) to be broken down by population subgroups (such as caste, religion, or region) and by dimensions (health, education, living standards). This feature is essential for identifying whether poverty reduction is driven by a decrease in the number of poor people (incidence) or a reduction in the intensity of their deprivation, a nuance often lost in unidimensional poverty measurements.

The global MPI structure constitutes of three dimensions including health, education and living standards and ten indicators with both dimensions and indicators equally weighted. A deprivation profile is established for each person pointing towards the deprived indicators. Along with it, a headcount ratio (H) indicating the proportion of multidimensionally poor people in the population and intensity of poverty (A) represented by the average percentage of weighted deprivations experienced by the poor is obtained. MPI is calculated as a multiplicative sum of $H \times A$ (NITI Aayog, 2023). The H and A are defined as :

$$H = \frac{1}{n} \sum_{i=1}^n \rho(g_i^0, w, k), A = \frac{1}{q} \sum_{i=1}^n (\rho(g_i^0, w, k) \times c_i), \text{ where } q = \sum_{i=1}^n \rho(g_i^0, w, k)$$

Consequently, the adjusted poverty headcount ratio (MPI) is calculated as :

$$MPI = M_0 = H \times A = \frac{1}{n} \sum_{i=1}^n (\rho(g_i^0, w, k) \times c_i)$$

The NITI Aayog constructed a national MPI adhering to the Alkire and Foster (AF) methodology employed in the global MPI creating an indigenized measure of assessing multidimensional poverty. While incorporating all ten global MPI indicators, the national index expands its scope to include maternal health and bank accounts, aligning with India's development priorities (NITI Aayog, 2023). India's Multidimensional Poverty Index (MPI) approach defines individuals under five years of age and within the working-age range of 15 to 54 for men and 15 to 49 for women as relevant population segments. However, when assessing child-adolescent mortality, a narrower age group is employed, including children aged 0 to 59 months and adolescents aged 10 to 18 years. In contrast to traditional monetary poverty assessments based on consumption expenditure, the national MPI provides a granular analysis of deprivations at the district level.

The dual cut off methodology employs majorly two steps namely identification and aggregation. Identification involves building a deprivation profile by applying cutoffs within an indicator and identifying the incidence of multidimensional poverty by applying a cut-off across all indicators. The deprivation of each individual is marked as either 0 or 1 indicating not deprived (0) and deprived (1). If the achievement of an individual i in indicator j is denoted by x_{ij} , the first order cut off for indicator j is denoted by z_j , and the status of the individual is denoted by g_{ij}^0 , then

$$g_{ij}^0 = 1 \text{ if } x_{ij} < z_j \text{ and } g_{ij}^0 = 0 \text{ otherwise for all } i = 1, 2, \dots, n \text{ and } j = 1, 2, \dots, d$$

The counting vector or deprivation score is the sum of the weighted status of all the indicators for an individual.

The deprivation score of the individual i can be denoted as:

$$c_i = w_1 g_{i1}^0 + w_2 g_{i2}^0 + \dots + w_j g_{ij}^0, \text{ or } c_i = \sum_{j=1}^d w_j g_{ij}^0$$

where c_i is the counting vector for individual i upto the j^{th} indicator, g_{ij}^0 is the status of each indicator and w_j is the weight assigned to each indicator.

The poverty cut-off also mentioned as the second order cut-off is deployed to be 0.33 (33%) is the minimum deprivation score as individuals having a deprivation score greater than or equal to the second order cut off are considered to be multidimensionally poor. A score less than second order cut off is replaced with 0.

Headcount Ratio (H), where q is the total number of multidimensionally poor people in a population n is computed as,

$$H = \frac{q}{n}$$

Similarly, the average proportion of deprivations experienced by multidimensionally poor individuals denoted by Intensity of Poverty (A) is computed as,

$$A = \frac{1}{q} \sum_{i=1}^q c_i(k)$$

MPI is represented as $M_0 = H \times A$,

$$\text{Or } H \times A = \frac{q}{n} * \frac{1}{q} \sum_{i=1}^q c_i(k) = \frac{1}{n} \sum_{i=1}^n c_i(k) = \frac{1}{n} \sum_{i=1}^n \sum_{j=1}^d w_j g_{ij}^{\circ}(k)$$

(where, $c_i(k)$ is the censured deprivation score up to the i^{th} individual and q is the number of multidimensionally poor individuals. Further, g_{ij} is household member status and w_j is the weight of j^{th} indicator. Here, n is the total population and d is the number of dimensions)

Table 1: Dimension, Indicators, and Derivation conditions (NITI Aayog)

Dimension of Poverty	Indicator	Deprivation condition	Weight
Health (1/3)	Nutrition	Any person under 70 years of age for whom nutritional information is available is undernourished ¹ .	1/6
	Child -Adolescent Mortality	A child under 18 has died in the household in the five years preceding the survey	1/12
	Maternal Health	Any woman in the household who has given birth in the 5 years preceding the survey, has not received at least 4 antenatal care visits for the most recent birth or has not received assistance from trained skilled medical personnel during the most recent childbirth	1/12
Education (1/3)	Years of Schooling	No eligible household member has completed six years of schooling	1/6
	School Attendance	Any school-aged child ³ is not attending school up to the age at which he/she would complete class 8	1/6
Standard of Living (1/3)	Cooking Fuel	A household cooks using solid fuel, such as dung, crops, shrubs, wood, charcoal, or coal	1/21
	Sanitation	The household has unimproved or no sanitation facility or it is improved but shared with other households	1/21
	Drinking Water	The household's source of drinking water is not safe or safe drinking water is a 30-minute walk or longer walk from home, roundtrip	1/21
	Electricity	The household has no electricity	1/21
	Housing	The household has inadequate housing materials in any of the three components: floor, roof, or wall	1/21
	Assets	The household does not own more than one of these assets: radio, TV, telephone, computer, animal cart, bicycle, motorbike, or refrigerator, and does not own a car or truck	1/21
	Bank Account	No household member has a bank account or a post office account.	1/21

$$\text{Deprivation Score} = \frac{1}{3}\text{Health}_i + \frac{1}{3}\text{Education}_i + \frac{1}{3}\text{Standard of Living}_i,$$

$$[\frac{1}{3}\text{Health}_i = \frac{1}{6}\text{Nutrition}_i + \frac{1}{12}\text{Child – Adolescent Mortality}_i + \frac{1}{12}\text{Maternal Health}_i$$

$$\frac{1}{3}\text{Education}_i = \frac{1}{6}\text{Years of schooling}_i + \frac{1}{6}\text{School Attendance}_i$$

$$\frac{1}{3}\text{Standard of Living}_i = \frac{1}{21}\text{Cooking Fuel}_i + \frac{1}{21}\text{Sanitation}_i + \frac{1}{21}\text{Drinking Water}_i + \frac{1}{21}\text{Electricity}_i + \frac{1}{21}\text{Housing}_i + \frac{1}{21}\text{Assets}_i + \frac{1}{21}\text{Bank Account}_i]$$

Data

The National Family Health Survey (NFHS), a representative of the Indian Demographic Health Survey (DHS), is executed by the International Institute for Population Sciences (IIPS), Mumbai under the mandate of India's Ministry of Health and Family Welfare. The primary objectives of the NFHS constitutes the provision of data and technical assistance to inform the formulation and implementation of health and family welfare policies and programs. Moreover, the NFHS offers valuable insights into contemporary health and family welfare challenges. This study employs micro survey data from three rounds of NFHS surveys, (NFHS-3, NFHS-4 and NFHS -5) analysing socio – economic characteristics such as age, caste, religion, place of residence, wealth status, household size etc of different social groups such as Scheduled Castes (SC), Scheduled Tribes (ST), Other Backward Classes (OBC) and Others (IIPS, n.d.).

Outcomes

Primarily, our research study presents a comprehensive analysis of multidimensional poverty across diverse socioeconomic and social groups in India, examining three critical time periods (2005-06, 2015-16, and 2019-21). Through the systematic computation of Multidimensional Poverty Index (MPI) values, we evaluated poverty dynamics across various socio-economic groups and geographical regions. Our investigation extended beyond conventional poverty metrics to capture the temporal evolution of multidimensional deprivation patterns across different social groups. The research systematically decomposed the contributing factors to MPI variations, with particular attention to the interplay between headcount ratios and intensity of poverty. Furthermore, we conducted an in-depth examination of specific deprivation indicators and their relative contributions to overall poverty

measures. By implementing decomposition techniques to analyze mean differences in MPI across groups, we identified and quantified the fundamental drivers of poverty disparities, thereby providing insights into both observed characteristics and underlying structural factors that perpetuate poverty across different social segments.

Socioeconomic Status Characteristics

The analysis incorporated multiple socioeconomic status indicators through a comprehensive set of demographic and social parameters. Age stratification comprised four distinct categories: early childhood (0–5 years), school-age (6–14 years), working-age adults (15–59 years), and elderly (60+ years), enabling examination across the complete lifespan. Marital status was classified into three categories: married, unmarried, and widowed/separated/divorced. Household composition was categorized by size: small (≤ 4 members), medium (5–6 members), and large (> 6 members). The analysis was deepened by examining the data through the lens of social groups, including caste (Scheduled Castes, Scheduled Tribes, Other Backward Classes, and Others) and religion (Hindu, Muslim, Christian, Sikh, and Others). Economic status was assessed through wealth quintiles (poorest, poorer, middle, richer, richest). Additional demographic variables comprising gender (male/female) and residential location (rural/urban) were also utilised in the analysis.

Statistical Analysis

Our analytical framework incorporates multiple methodological approaches to examine multidimensional poverty across social groups. Initially, we compute the Multidimensional Poverty Index (MPI) values across various socioeconomic status (SES) categories by calculating their respective headcount ratios and intensities spanning three National Family Health Survey (NFHS) periods (2005–06, 2015–16, and 2019–21). This analysis is extended to state-level disaggregation of MPI values across different social groups for the corresponding time periods. To assess the probability of experiencing multidimensional poverty by background characteristics, we employ pooled logistic regression analysis, generating odds ratios based on unadjusted headcount ratios while controlling for relevant covariates performed using STATA 17 and MLwiN 3.09. We further employ the Kitagawa Decomposition Method to analyze the relative contributions of headcount ratio and intensity to multidimensional poverty across social groups over consecutive NFHS periods (NFHS 3–4, NFHS 4–5, and NFHS 3–5), along with the net differences in MPI across these intervals. Finally, we implement the Blinder–Oaxaca Decomposition to analyse mean differences in multidimensional poverty across different groups over the appended time periods (NFHS 3–4, NFHS 4–5, and NFHS 3–5). To ensure the robustness of the trend analysis, the comparability of indicators across the three NFHS rounds

was carefully maintained. While minor changes in survey questionnaires occurred over the 15-year period, the definitions of the core 12 indicators used for the MPI construction—specifically regarding nutrition, mortality, and standard of living assets—were harmonized to match the standardized definitions utilized by the Oxford Poverty and Human Development Initiative (OPHI) and NITI Aayog for longitudinal analysis. Finally, it is imperative to note that the factors identified through the logistic regression and decomposition analyses represent statistical associations. Given the cross-sectional nature of the NFHS data, these findings should be interpreted as correlational rather than establishing definitive causal relationships between socioeconomic characteristics and multidimensional poverty.

Decomposition

• Kitagawa Decomposition

Decomposition techniques serve as essential analytical tools for disaggregating demographic variables into their constituent components, thereby addressing the methodological challenges posed by confounding compositional effects (Canudas Romo, 2003). Building upon Kitagawa's (1955) seminal work on 'components of a difference between two rates,' which advanced beyond the limitations of traditional standardisation techniques, our analytical framework focuses on decomposing inequalities across distinct social groups (Scheduled Castes, Scheduled Tribes, Other Backward Classes, and Others). The temporal analysis spans three intervals of the National Family Health Survey (NFHS): between NFHS-3 and NFHS-4, NFHS-4 and NFHS-5, and the comprehensive period between NFHS-3 and NFHS-5. This approach enables the examination of variations in both the headcount ratio (H) and intensity (A) of multidimensional poverty across different socio-economic strata.

The overall rate of difference in headcount ratio across two groups can be decomposed as:

$$\Delta = MPI^A - MPI^B = \sum_i H_i^B \cdot A_i^B - \sum_i H_i^A \cdot A_i^A$$

where MPI denotes the value of Multidimensional Poverty Index, H denotes headcount ratio and A represents the intensity of poverty across two time periods A and B.

Each of these terms will be divided into two equal parts with certain additional terms added and subtracted, implying that the difference (Δ) will be constant:

$$\Delta = \frac{\sum_i H_i^B \cdot A_i^B}{2} + \frac{\sum_i H_i^B \cdot A_i^B}{2} - \frac{\sum_i H_i^A \cdot A_i^A}{2} - \frac{\sum_i H_i^A \cdot A_i^A}{2} + \frac{\sum_i H_i^B \cdot A_i^A}{2} - \frac{\sum_i H_i^B \cdot A_i^A}{2} + \frac{\sum_i H_i^A \cdot A_i^B}{2} - \frac{\sum_i H_i^A \cdot A_i^B}{2}$$

Therefore,

$$\Delta = \sum_i (H_i^B - H_i^A) \cdot \left[\frac{A_i^B + A_i^A}{2} \right] + \sum_i (A_i^B - A_i^A) \cdot \left[\frac{H_i^B + H_i^A}{2} \right].$$

Here, the proportion attributable to each category is calculated over the specific decomposition time periods (Preston et al., 2001).

Oaxaca – Blinder Decomposition

The Oaxaca – Blinder decomposition facilitates an extension of the Kitagawa Decomposition method as a counterfactual decomposition technique applied in terms of linear regression models. Primarily, it sets forth the estimation of mean difference in the outcome variable between two different groups. This is further supplemented with the fact that the OB decomposition emphasizes on the interaction between the group parameter differences as well as the group characteristics (Oaxaca, 1973; Blinder, 1973).

Within this analytical framework, we designate two population groups, A and B, corresponding to the respective time periods in the National Family Health Survey (NFHS) rounds (3, 4, and 5). The outcome variable Y , designated for decomposition, represents the aggregate deprivations experienced by an individual, computed as the product of the unadjusted headcount ratio and the counting vector. The sample sizes for populations A and B are denoted by N_A and N_B respectively. The methodological approach incorporates a set of K mutually exclusive indicator variables, consolidated into a single categorical explanatory variable X , such that

$$\sum_{k=1}^K X_{ik} \forall i$$

where i denotes the i th individual within a sample group. The outcome indicator variables for the respective population groups are represented by Y_i^A and Y_i^B (Oaxaca & Sierminska, 2025).

The group mean proportions are given by

$$\bar{Y}^A = \frac{\sum_{i=1}^{N^A} Y_i^A}{N^A} = \frac{N_Y^A}{N^A}$$

$$\bar{Y}^B = \frac{\sum_{i=1}^{N^B} Y_i^B}{N^B} = \frac{N_Y^B}{N^B}$$

Hence,

$$\bar{Y}^A - \bar{Y}^B = \frac{N_Y^A}{N^A} - \frac{N_Y^B}{N^B}$$

Let the outcome rates in the k th category of the predictor variable X for groups A and B be:

$$\begin{aligned}\bar{Y}_k^A &= \frac{\sum_{i=1}^{N^A} Y_i^A X_{ik}}{N_k^A}, k = 1, \dots, K \\ &= \frac{N_{Yk}^A}{N_k^A} \\ \bar{Y}_k^B &= \frac{\sum_{i=1}^{N^B} Y_i^B X_{ik}}{N_k^B}, k = 1, \dots, K \\ &= \frac{N_{Yk}^B}{N_k^B}\end{aligned}$$

The linear probability regression model for the two population groups is presented as:

$$\begin{aligned}Y_i^A &= \sum_{k=1}^K X_{ik} \beta_k^A + \varepsilon_i, i = 1, \dots, N^A \\ Y_i^B &= \sum_{k=1}^K X_{ik} \beta_k^B + \varepsilon_i, i = 1, \dots, N^B\end{aligned}$$

The OLS estimator corresponding to the separate linear regressions of Y_i on each X_{ik} indicator variable will be :

$$\begin{aligned}b_k^A &= \frac{\sum_{i=1}^{N^A} X_{ik} Y_i^A}{\sum_{i=1}^{N^A} X_{ik}^2}, k = 1, \dots, K \\ &= \frac{N_{Yk}^A}{N_k^A} = \bar{Y}_k^A \\ b_k^B &= \frac{\sum_{i=1}^{N^B} X_{ik} Y_i^B}{\sum_{i=1}^{N^B} X_{ik}^2}, k = 1, \dots, K \\ &= \frac{N_{Yk}^B}{N_k^B} = \bar{Y}_k^B\end{aligned}$$

In Blinder Oaxaca Decomposition, one group is assigned as the reference group and we execute the simple averages of the two groups estimated coefficients such that,

$$b_k^* = \frac{(b_k^A + b_k^B)}{2}, k = 1, \dots, K = \left(\frac{\bar{Y}_k^A + \bar{Y}_k^B}{2} \right)$$

Therefore, the OLS decomposition can be explained as the following :

$$\begin{aligned}\bar{Y}^A - \bar{Y}^B &= \sum_{k=1}^K \bar{X}_k^A b_k^A - \sum_{k=1}^K \bar{X}_k^B b_k^B \\ &= \sum_{k=1}^K (\bar{X}_k^A - \bar{X}_k^B) b_k^* + \sum_{k=1}^K \bar{X}_k^A (b_k^A - b_k^*) + \sum_{k=1}^K \bar{X}_k^B (b_k^* - b_k^B)\end{aligned}$$

The initial component of the equation, represented by

$$\sum_{k=1}^K (\bar{X}_k^A - \bar{X}_k^B) b_k^*$$

quantifies the differential attributed to group variations in the explanatory variables, termed as the endowment effect . Along with it, the second component

$$\sum_{k=1}^K \bar{X}_k^A (b_k^A - b_k^*)$$

comprises the contribution arising from differences in coefficients, constituting the unexplained component, designated as the coefficients effect. Furthermore, interaction effect which is represented by the final part of the equation

$$\sum_{k=1}^K \bar{X}_k^B (b_k^* - b_k^B)$$

captures the simultaneous manifestation of differences in both endowments and coefficients between the two groups under consideration (Jann, 2008; Rahimi & Hashemi Nazari, 2021).

Results

India's MPI exhibited significant reductions across three NFHS rounds (2005–06 to 2019–21), declining substantially from 0.282 to 0.066 (Table 2.1). This decline was driven by both a decrease in the headcount ratio (from 52.3% to 14.7%) and a moderate improvement in intensity (from 53.9% to 44.5%). The annual rate of reduction increased from 8.5% between NFHS 3 and 4 to 10.5% between NFHS 4 and 5. Among demographic groups, changes were observed in wealth quintiles, with the poorest wealth quintile experiencing a reduction in multidimensionally poor population, declining from 92.8% to 42.1% between NFHS 3 and 5. While children aged 0–14 years demonstrated improvements (from 63.0% to 23.4%), they consistently remained the most vulnerable age group across all rounds. The Scheduled Tribes consistently exhibited the highest poverty rates (26.8% in 2019–21), significantly exceeding other groups. The rural–urban divide persisted, with rural areas (18.9%) exhibiting higher multidimensionally poor populations compared to urban areas (5.3%) in 2019–21. Furthermore, despite similar household characteristics, females consistently displayed slightly higher MPI scores than males across all three rounds. (Table 2.2)

Table 2.1 : Multidimensional Poverty Headcount(H), Intensity(A), and Index values by background characteristics, India

Background Variables	NFHS-3 (2005-06)			NFHS-4 (2015-16)			NFHS-5 (2019-21)		
	H	A	MPI	H	A	MPI	H	A	MPI
Age									
0-14 yrs.	0.630	0.571	0.360	0.348	0.494	0.172	0.234	0.462	0.108
15-59 yrs.	0.466	0.522	0.243	0.200	0.461	0.092	0.116	0.436	0.051
60+ yrs.	0.500	0.490	0.245	0.217	0.436	0.094	0.115	0.413	0.048
Sex									
Male	0.512	0.537	0.275	0.232	0.470	0.109	0.139	0.444	0.062
Female	0.535	0.541	0.289	0.255	0.473	0.121	0.155	0.446	0.069
Household size									
<=4 members	0.437	0.502	0.22	0.173	0.453	0.078	0.097	0.432	0.042
5 to 6 members	0.515	0.547	0.281	0.234	0.481	0.113	0.148	0.455	0.067
6+ members	0.620	0.558	0.346	0.349	0.476	0.166	0.228	0.444	0.101
Marital Status									
Married	0.496	0.526	0.261	0.220	0.462	0.102	0.130	0.435	0.057
Unmarried	0.453	0.524	0.238	0.151	0.449	0.068	0.078	0.430	0.033
Widow/separated/ Divorced	0.535	0.506	0.271	0.237	0.446	0.106	0.129	0.423	0.055
Wealth Quintile									
Richest	0.055	0.404	0.022	0.012	0.376	0.004	0.010	0.380	0.004
Richer	0.259	0.440	0.114	0.051	0.398	0.020	0.033	0.390	0.013
Middle	0.596	0.477	0.284	0.148	0.413	0.061	0.077	0.403	0.031
Poorest	0.825	0.539	0.445	0.337	0.445	0.150	0.185	0.420	0.078
Poorer	0.928	0.619	0.575	0.654	0.506	0.331	0.421	0.469	0.198
Caste									
SC	0.632	0.545	0.345	0.286	0.474	0.136	0.182	0.450	0.082
ST	0.778	0.587	0.457	0.436	0.490	0.214	0.268	0.458	0.123
OBC	0.552	0.532	0.294	0.240	0.468	0.112	0.138	0.439	0.061
Others	0.341	0.514	0.176	0.137	0.457	0.063	0.074	0.433	0.032
Religion									
Hindu	0.524	0.532	0.279	0.242	0.467	0.113	0.144	0.441	0.063
Muslim	0.590	0.583	0.344	0.297	0.498	0.148	0.189	0.46	0.087
Christian	0.367	0.513	0.188	0.140	0.463	0.065	0.101	0.453	0.046
Sikh	0.218	0.473	0.103	0.058	0.433	0.025	0.048	0.419	0.020
Others	0.477	0.529	0.252	0.211	0.472	0.099	0.140	0.446	0.062
Place of Residence									
Rural	0.651	0.547	0.356	0.319	0.474	0.151	0.189	0.446	0.085
Urban	0.243	0.492	0.120	0.084	0.454	0.038	0.053	0.432	0.023
All India	0.523	0.539	0.282	0.243	0.472	0.115	0.147	0.445	0.066

Table 2.2: Multidimensional Poverty Index for Social Groups, Indian States (NFHS-5)

State /UT	SC			ST			OBC			Others			All India	
	H	A	MPI	H	A	MPI	H	A	MPI	H	A	MPI	MPI	MPI
A & N Island	0.078	0.345	0.027	0.036	0.403	0.014	0.003	0.413	0.001	0.032	0.41	0.013	0.01	0.01
Andhra Pradesh	0.076	0.411	0.031	0.256	0.433	0.111	0.062	0.404	0.025	0.018	0.371	0.007	0.026	0.026
Arunachal Pradesh	0.127	0.427	0.054	0.129	0.419	0.054	0.219	0.475	0.104	0.168	0.456	0.076	0.06	0.06
Assam	0.164	0.431	0.071	0.126	0.432	0.054	0.159	0.451	0.072	0.173	0.439	0.076	0.086	0.086
Bihar	0.458	0.492	0.225	0.466	0.502	0.234	0.308	0.462	0.142	0.197	0.463	0.091	0.158	0.158
Chandigarh	0.03	0.381	0.011	-	-	-	0.06	0.433	0.026	0.025	0.586	0.015	0.017	0.017
Chhattisgarh	0.143	0.416	0.059	0.268	0.444	0.119	0.109	0.41	0.045	0.041	0.391	0.016	0.067	0.067
D & NH	0.023	0.436	0.01	0.169	0.414	0.07	0.048	0.421	0.02	0.041	0.463	0.019	0.042	0.042
Goa	0	.	0	0.016	0.346	0.006	0.001	0.357	0	0.004	0.341	0.001	0.003	0.003
Gujarat	0.109	0.418	0.045	0.256	0.451	0.116	0.111	0.425	0.047	0.031	0.403	0.013	0.051	0.051
Haryana	0.101	0.419	0.042	0.094	0.428	0.04	0.102	0.449	0.046	0.027	0.426	0.012	0.032	0.032
Himachal Pradesh	0.089	0.393	0.035	0.077	0.392	0.03	0.022	0.4	0.009	0.037	0.408	0.015	0.02	0.02
Jammu & Kashmir	0.061	0.417	0.026	0.145	0.439	0.064	0.091	0.422	0.038	0.03	0.413	0.012	0.021	0.021
Jharkhand	0.35	0.46	0.161	0.384	0.479	0.184	0.216	0.434	0.094	0.13	0.436	0.057	0.128	0.128
Karnataka	0.098	0.412	0.04	0.133	0.436	0.058	0.062	0.405	0.025	0.049	0.395	0.02	0.032	0.032
Kerala	0.012	0.374	0.005	0.066	0.405	0.027	0.006	0.356	0.002	0.001	0.352	0	0.002	0.002
Ladakh	0	.	0	0.046	0.408	0.019	-	-	-	0.008	0.405	0.003	0.017	0.017
Lakshadweep	0	.	0	0.009	0.376	0.003	0	.	0	0	.	0	0.003	0.003
Madhya Pradesh	0.225	0.434	0.098	0.357	0.457	0.163	0.157	0.423	0.066	0.076	0.409	0.031	0.087	0.087
Maharashtra	0.067	0.405	0.027	0.243	0.445	0.108	0.047	0.393	0.019	0.047	0.405	0.019	0.032	0.032
Manipur	0.116	0.429	0.05	0.155	0.424	0.066	0.062	0.399	0.025	0.047	0.404	0.019	0.036	0.036
Meghalaya	0.033	0.425	0.014	0.302	0.479	0.145	0.171	0.589	0.101	0.199	0.468	0.093	0.136	0.136
Mizoram	0.189	0.469	0.089	0.051	0.451	0.023	0.333	0.478	0.159	-	-	-	0.026	0.026
Nagaland	0.133	0.429	0.057	0.157	0.424	0.066	0.041	0.421	0.017	0.223	0.423	0.095	0.066	0.066

Table 2.2: Multidimensional Poverty Index for Social Groups, Indian States (NFHS-5)

State /UT	SC			ST			OBC			Others			All India	
	H	A	MPI	H	A	MPI	H	A	MPI	H	A	MPI	MPI	MPI
NCT of Delhi	0.047	0.423	0.02	0.039	0.51	0.02	0.03	0.425	0.013	0.026	0.415	0.011	0.015	0.015
Odisha	0.15	0.429	0.064	0.328	0.468	0.154	0.078	0.419	0.033	0.038	0.4	0.015	0.067	0.067
Puducherry	0.01	0.393	0.004	0	.	0	0.009	0.376	0.003	0	.	0	0.003	0.003
Punjab	0.077	0.414	0.032	0.064	0.399	0.025	0.03	0.412	0.013	0.016	0.39	0.006	0.02	0.02
Rajasthan	0.19	0.432	0.082	0.262	0.446	0.117	0.13	0.419	0.054	0.074	0.411	0.03	0.065	0.065
Sikkim	0.057	0.43	0.025	0.035	0.39	0.014	0.021	0.431	0.009	0	.	0	0.012	0.012
Tamil Nadu	0.036	0.384	0.014	0.065	0.425	0.028	0.018	0.383	0.007	0.007	0.419	0.003	0.009	0.009
Telangana	0.065	0.408	0.027	0.124	0.425	0.053	0.05	0.402	0.02	0.022	0.393	0.009	0.024	0.024
Tripura	0.121	0.41	0.05	0.205	0.443	0.091	0.066	0.411	0.027	0.076	0.42	0.032	0.056	0.056
Uttar Pradesh	0.287	0.451	0.13	0.353	0.484	0.171	0.227	0.446	0.101	0.129	0.445	0.058	0.101	0.101
Uttarakhand	0.119	0.426	0.051	0.083	0.407	0.034	0.149	0.423	0.063	0.052	0.401	0.021	0.039	0.039
West Bengal	0.123	0.421	0.052	0.235	0.447	0.105	0.075	0.417	0.031	0.094	0.43	0.041	0.05	0.05
All India	0.182	0.449	0.082	0.269	0.458	0.123	0.138	0.439	0.061	0.075	0.433	0.032	0.066	0.066

H: Headcount Ratio, A: Intensity, MPI: Multidimensional Poverty Index Scores

The MPI for Indian states reveals significant disparities in poverty levels across different social groups in India (Table 3). Examining the NFHS-5 (2019-21) data, the Scheduled Tribe population emerges as the most disadvantaged population group with several states showing high headcount ratios and MPI scores. Jharkhand (H: 54.6%, MPI: 0.271), Chhattisgarh (H: 42.1%, MPI: 0.185), and Madhya Pradesh (H: 57.1%, MPI: 0.280) have the highest poverty levels among the ST population. The Scheduled Caste and Other Backward Classes also exhibit elevated poverty, though generally lower than the STs. Comparing the NFHS-4 (2015-16) and NFHS-3 (2005-06) data, some states have shown progress in reducing multidimensional poverty, particularly among the ST population. For instance, Andhra Pradesh witnessed a significant decrease in ST headcount ratio (from 73% to 40.2%) and MPI (from 0.420 to 0.181) over this period. Rajasthan also saw a decline in ST headcount ratio (from 89.1% to 58.8%) and MPI (from 0.532 to 0.304). However, other states like Uttar Pradesh and Jharkhand continue to struggle with high poverty levels, especially among the ST and SC communities, with limited improvement over the years. Notably, the intensity values, remain relatively similar across social groups and states.

A pooled regression analysis examining the multidimensional poverty headcount ratio revealed that Scheduled Castes (NFHS-5: OR 1.62, 95% CI: 1.59-1.65), Scheduled Tribes (NFHS-5: OR 1.75, CI: 1.71-1.78), and Other Backward Classes (NFHS-5: OR 1.42, CI: 1.4-1.44) consistently demonstrated significantly higher likelihood of being multidimensionally poor across all three rounds (2005-06 to 2019-21) (Table 4). The poorest quantile showed a substantial reduction in the percentage of population experiencing multidimensional poverty, decreasing from 203.17 in NFHS-3 to 76.82 in NFHS-5. Similarly, the middle and poorer wealth quintiles showed significant decrease in odds of being multidimensional poor over the three rounds while being relatively higher in the same round than the richest and richer wealth quintiles. Households with more than six members exhibited increased likelihood of being multidimensionally poor (NFHS-5: OR 3.8, 95% CI: 3.75-3.84), while the 15-59 years' age cohort demonstrated a lower probability of multidimensional poverty (NFHS-5: OR 0.79, CI: 0.77-0.81) than under 14 age group. Rural populations, and certain religious groups, like Hindus and Muslims, also faced significantly higher risks of being multidimensionally poor. Despite similar household characteristics, females showed slightly more odds of being multidimensionally poor than males across all three rounds (NFHS-5: OR 1.07, CI: 1.06-1.08).

The Kitagawa Decomposition analysis of multidimensional poverty headcount ratio and intensity across three NFHS survey periods was conducted (Table 5). The maximum net reduction in the MPI was observed between the NFHS 3 and NFHS 5 (0.216). The net MPI reductions observed between the consecutive survey rounds of NFHS 3-4 and NFHS 4-5 were 0.167 and 0.049, respectively. The headcount ratio

Table 3: Pooled logistic Regression based odds ratio (OR) for unadjusted headcount (H) by background characteristics

Background Characteristics	NFHS-3	NFHS-4	NFHS-5
	OR [95% CI]	OR [95% CI]	OR [95% CI]
Age			
0-14 yrs. (ref)	1	1	1
15-59 yrs.	0.69*** (0.66, 0.71)	0.84*** (0.83, 0.86)	0.79*** (0.77, 0.81)
60+ yrs.	0.59*** (0.56, 0.62)	0.77*** (0.75, 0.78)	0.68*** (0.66, 0.70)
Sex			
Male (ref)	1	1	1
Female	1.06*** (1.04, 1.08)	1.08*** (1.07, 1.09)	1.07*** (1.06, 1.08)
Household Size			
Household size <= 4 (ref)	1	1	1
Household size 5-6	1.39*** (1.36, 1.42)	1.33* (1.31, 1.34)	1.55*** (1.53, 1.56)
Household size >6	3.28*** (3.21, 3.36)	3.27*** (3.24, 3.31)	3.80*** (3.75, 3.84)
Marital Status			
Married	1.75*** (1.70, 1.79)	2.16*** (2.13, 2.18)	2.33*** (2.30, 2.36)
Unmarried (ref)	1	1	1
Widow/Seperated/Divorced	1.81*** (1.73, 1.89)	2.29*** (2.25, 2.34)	
Wealth Quintile			
Richest (ref)	1	1	1
Richer	5.50*** (5.29, 5.73)	4.09*** (3.95, 4.24)	3.17*** (3.05, 3.30)
Middle	25.12*** (24.13, 26.15)	12.75*** (12.33, 13.19)	7.89*** (7.60, 8.21)
Poorer	78.01*** (74.70, 81.46)	38.64*** (37.36, 39.95)	21.21*** (20.43, 22.03)
Poorest	203.17*** (193.44, 213.40)	153.29*** (148.19, 158.56)	76.82*** (73.97, 79.79)
Caste			
SC	1.57*** (1.52, 1.61)	1.37*** (1.35, 1.39)	1.62*** (1.59, 1.65)
ST	1.83*** (1.76, 1.91)	1.71*** (1.69, 1.74)	1.75*** (1.71, 1.78)
OBC	1.38*** (1.35, 1.41)	1.36*** (1.34, 1.38)	1.42*** (1.40, 1.44)
Others (ref)	1	1	1
Religion			
Hindu	1.15*** (1.09, 1.22)	1.12*** (1.08, 1.15)	0.98* (0.94, 1.01)
Muslim	2.21*** (2.07, 2.35)	2.16*** (2.10, 2.24)	1.84*** (1.77, 1.92)
Christian	0.97* (0.89, 1.05)	0.81*** (0.77, 0.85)	0.75*** (0.71, 0.79)
Others (ref)	1	1	1
Area			
Rural	1.26*** (1.23, 1.28)	1.14*** (1.13, 1.16)	1.02*** (1.01, 1.04)
Urban (ref)	1	1	1

demonstrated a substantial contribution to the observed MPI reduction throughout all survey rounds, increasing from 85% to 89% across consecutive periods (NFHS 3-4 and NFHS 4-5). The most significant MPI reduction was reported among the ST population, which experienced a 0.334 reduction between NFHS 3 and NFHS 5. This substantial improvement can be attributed to both a decrease in the proportion of the ST population living in multidimensional poverty and a decrease in the intensity of their poverty. In contrast, the proportion attributable to intensity marginally decreased across other social groups.

The analysis of the contribution of various indicators across the NFHS periods reveals that the Standard of Living dimension had the most significant contribution to the overall MPI in NFHS 3 (Table 5). However, this trend shifted over time, with the Health dimension emerging as the largest contributor, accounting for 42.8% of the MPI in NFHS 5. The Nutrition indicator was consistently the most significant contributor across all three NFHS rounds, demonstrating almost 6% increase in its contribution, from NFHS 3 to NFHS 5. This was followed by the education dimension, with the Years of Schooling indicator showing an increase from 13% in NFHS 3 to 17% in NFHS 5 accompanied by a decreasing headcount ratio. Within the Standard of Living dimension, the Bank Account indicator saw a significant decline, decreasing from a 7% contribution in NFHS 3 to just 1% in NFHS 5. While, certain other indicators, such as Assets, and Electricity, demonstrated substantial reductions in their contributions, indicators such as Housing and Cooking Fuel, maintained relatively consistent contributions.

The Blinder-Oaxaca decomposition analysis of the changes in the MPI across the NFHS periods was conducted (Table 6). Examining the household size, the endowment effect shows a positive contribution of 1.32% between NFHS 3-4 and 3.04% between NFHS 4-5, indicating that that population shift towards smaller household sizes has been potentially favourable in reducing multidimensional poverty. The positive coefficient effect for larger household sizes is 6.93% and 9.04%, respectively, suggesting that the adverse impact of larger household sizes on multidimensional poverty has decreased over time. The endowment effects for the Poorest and Poorer wealth groups are substantial and negative, at -3.11% and -1.48% respectively, implying that the population movements towards these economically disadvantaged groups have hindered progress in alleviating multidimensional poverty. However, the coefficient effects for the Poorest (87.25%), Poorer (94.52%), and Middle (38.54%) wealth quintiles are positive and significant, indicating that the relationship between poverty and these wealth groups has become less severe. The analysis of social groups shows that the endowment effects for Scheduled Castes (-0.54%), Scheduled Tribes (-0.42%), and Other Backward Classes (-0.18%) are negative, suggesting that the demographic shifts towards these socially marginalized communities have contributed to the persistence of multidimensional deprivation. The coefficient effects for these social

Table 4: Kitagawa Decomposition for Contribution of Headcount(H) & Intensity(A) to Multidimensional Poverty Index

Caste Groups	MPI Difference			MPI Difference			MPI Difference				
	NFHS 3 – NFHS 4			NFHS 4 – NFHS 5			NFHS 3 – NFHS 5				
	Prop.	Att.	To H	Prop.	Att.	To H	Prop.	Att.	To H		
SC	84%	16%		0.209	89%		11%	0.054	85%	15%	0.263
ST	76%	24%		0.243	88%		12%	0.091	80%	20%	0.334
OBC	86%	14%		0.182	90%		10%	0.052	86%	14%	0.233
Others	88%	12%		0.113	92%		8%	0.03	88%	12%	0.143
All India	85%	15%		0.167	89%		11%	0.049	85%	15%	0.216

Table 5: Contribution of deprivations to Multidimensional Poverty Index Headcount Values

Dimensions	Indicators	Adjusted HCR			Indicator Contribution			Dimension Contribution		
		NFHS-3	NFHS-4	NFHS-5	NFHS-3	NFHS-4	NFHS-5	NFHS-3	NFHS-4	NFHS-5
Education	Years of schooling	0.04	0.02	0.01	13.30%	15.50%	16.80%			
	School Attendance	0.03	0.01	0.01	11.10%	7.60%	9.20%	24.30%	23.00%	26.00%
Health	Nutrition	0.07	0.03	0.02	24.40%	28.00%	29.80%			
	Child-Adolescent Mortality	0	0	0	1.00%	1.30%	1.50%	32.30%	39.20%	42.80%
Standard of Living	Maternal Health	0.02	0.01	0.01	6.90%	9.90%	11.50%			
	Electricity	0.01	0	0	4.60%	3.40%	1.30%			
	Sanitation	0.02	0.01	0	8.00%	8.60%	6.60%			
	Drinking Water	0.01	0	0	2.60%	2.10%	1.60%			
	Housing	0.02	0.01	0.01	7.10%	8.40%	8.60%	43.40%	37.70%	31.20%
	Cooking Fuel	0.02	0.01	0.01	8.40%	9.40%	8.80%			
	Bank Account	0.02	0	0	6.60%	2.20%	0.80%			
	Assets	0.02	0	0	6.00%	3.70%	3.40%			
Total	0.282	0.115	0.066	100.00%	100.00%	100.00%	100.00%	100.00%	100.00%	

Table 6: Blinder–Oaxaca Decomposition of the Multidimensional Poverty Index by Background Characteristics: Contribution to Total Effects

	NFHS 3–4	NFHS 3–5	NFHS 4–5
Endowment Effect			
Age	1.26%	1.25%	1.62%
Sex	0.06%	0.00%	0.00%
Household Size	1.32%	1.52%	3.04%
Marital Status	0.06%	0.14%	0.00%
WQ	–3.11%	–1.48%	–0.20%
Caste	–0.54%	–0.42%	–0.41%
Religion	–0.18%	0.00%	0.41%
Area	0.00%	0.00%	0.20%
States/UTs	–1.50%	–5.08%	–27.59%
Coefficient effect			
Age	–5.03%	–5.54%	–6.49%
Sex	4.91%	–0.79%	–2.23%
Household Size	9.04%	6.93%	2.03%
Marital Status	–1.38%	–1.06%	0.00%
WQ	84.91%	87.25%	94.52%
Caste	8.14%	6.93%	2.43%
Religion	2.40%	2.91%	5.07%
Area	4.73%	4.20%	2.03%
States/UTs	17.07%	16.95%	38.54%
Cons.	–22.04%	–28.45%	–67.95%
Interaction effect			
Age	0.42%	0.60%	0.20%
Sex	–0.12%	0.00%	0.00%
Household Size	0.72%	1.34%	0.41%
Marital Status	–0.30%	–0.28%	0.00%
WQ	–2.28%	–2.91%	0.00%
Caste	–0.66%	–0.74%	0.00%
Religion	–0.12%	0.00%	0.20%
Area	0.06%	–0.05%	0.00%
States/UTs	1.92%	16.49%	54.36%
Total Effect	0.167	0.217	0.049

groups are also positive, implying that the inherent disadvantages associated with these communities have diminished.

Examining regional disparities, the endowment effect for states such as Bihar (–1.50%), Odisha (–1.50%), and Jharkhand (–5.08%) is negative, reflecting that the concentration of vulnerable populations within these regions continues to pose a challenge to overall poverty reduction. Conversely, the coefficient effects for Uttar

Pradesh (38.54%), Bihar (38.54%), West Bengal (38.54%), and Andhra Pradesh (4.73%) are positive and significant. This suggests that state-specific structural factors and regional characteristics, such as governance, infrastructure, and state-level policy implementation, have become less associated with multidimensional poverty over time, indicating a relative improvement in the environment for poverty reduction in these states.

Discussion

The analysis of multidimensional poverty across three NFHS rounds revealed several key insights. Firstly, the Multidimensional Poverty Index demonstrated a significant decline across the three NFHS rounds, with the most substantial reductions observed among the poorest wealth quintiles, rural populations, and children aged 0–14 years. Notably, children, Scheduled Tribes, rural populations, and females consistently exhibited the highest poverty rates, with states like Uttar Pradesh and Jharkhand continuing to manifest multidimensional poverty. Additionally, despite overall national-level improvements, the Scheduled Tribe population's multidimensional poverty remained relatively high. Furthermore, the health dimension, particularly the nutrition indicator, emerged as the most significant contributor to MPI, accounting for the largest share of multidimensional poverty in 2019–21. The reductions in MPI were primarily attributable to improvements in weighted headcount ratios, with notable advancements in publicly provided variables such as health, education (years of schooling), bank accounts, assets, and electricity access. Lastly, the Blinder–Oaxaca decomposition revealed that, while population movements towards economically disadvantaged groups appeared to hinder poverty reduction, the analysis showed some improvements. The coefficient effects were positive and significant for the Poorest, Poorer, and Middle wealth quintiles, indicating a diminishing severity of poverty. Conversely, for Scheduled Castes, Scheduled Tribes, and Other Backward Classes, negative endowment effects suggested persistent demographic challenges, even as the coefficient effects implied reducing inherent disadvantages across different social groups and states.

The analysis reveals a significant reduction in multidimensional poverty in India, declining to 0.066 by 2019–21, aligning with NITI Aayog's findings (NITI Aayog, 2023). The national headcount ratio demonstrated a substantial 38% reduction between 2005–06 and 2019–21, effectively challenging South Asia's historical narrative of widespread poverty (OPHI, 2018). This reduction was most pronounced among the poorest wealth quintiles, with vulnerable populations experiencing more substantial improvements in absolute terms (Alkire & Seth, 2015; Alkire et al., 2021). Geographically, the poverty landscape revealed significant disparities, with states like Bihar, Meghalaya, Jharkhand, and Uttar Pradesh consistently demonstrating the highest Multidimensional Poverty Index (MPI) levels (Tripathi &

Yenneti, 2020). Furthermore, the poverty indicators remained significantly higher in rural areas compared to urban regions (Das et al., 2021). Additionally, women consistently experienced a higher likelihood of being multidimensionally poor. This trend extends beyond India, as evidenced by research on African nations, where women face similar challenges due to factors such as limited access to education, healthcare, and economic opportunities, as well as social norms and cultural practices that perpetuate gender inequality (Rogan, 2016).

The percentage of children living in severe child food poverty accounts for 40% in India (UNICEF, 2024). Furthermore, health emerged as the most critical dimension, contributing over 40% to the MPI, with nutrition alone accounting for nearly 30% of multidimensional poverty. The Global Hunger Index's ranking of India at 105th further highlights the persistent challenges of the 'Triple Burden of Malnutrition' (Undernutrition, Overnutrition, and Micronutrient Deficiency) (UNICEF, 2024; WHH et al., 2024). Approximately 40% of children experience severe food poverty, with alarming rates of stunting (35.5%), wasting (19.3%), and underweight (32.1%) conditions (NFHS, 2021). Low-quality infant and young child feeding (IYCF) practices significantly contribute to poor nutritional outcomes (Menon et al., 2015; Ramachandran, 2010), while the burden of nutritional deprivation disproportionately affects vulnerable groups, including poor households, scheduled castes, children of underweight mothers, and illiterate women compared to affluent groups (Prasad et al., 2021). Raghunathan et al. (2021) estimated that 63–76% of rural poor cannot afford recommended diets, highlighting the critical need for targeted nutrition interventions and safety net programs. Studies have consistently emphasized that affordable prices coupled with safety net programs covering nutritional norms, dietary allowance, and adjusted calorie intake are vital in mitigating the multidimensional aspects of nutrition deprivation (Raghunathan et al., 2021; Ryckman et al., 2021; Bhuyan et al., 2020).

Consistent with the existing literature, our analysis revealed that the Scheduled Tribe population demonstrated the highest Multidimensional poverty index values (Alkire et al., 2021; Pradhan et al., 2022). The rate of multidimensional poverty among Scheduled Tribes and Scheduled Castes significantly exceeds other socioeconomic groups in India, stemming from complex historical and contemporary social dynamics characterized by systemic discrimination and social injustice (Sahoo et al., 2023; Rupavath, 2023). Scheduled Tribes face a compounded disadvantage of socio-economic and spatial marginalization, with job insecurity in the unorganized sector being a substantial contributor to their economic vulnerability (Singh, 2019). The limited participation in economic growth among STs substantially widens the inequality scale, with chronic poverty largely attributable to social exclusion that systematically denies equitable resource access (Dubey, 2009; Mehta & Shah, 2003). Empirical studies, including our findings and research by Kaibarta et al. (2022) and Pradhan et al. (2022), consistently demonstrate that the standard of living

dimension contributes most significantly to the ST population's multidimensional poverty. While the Government of India has established the Ministry of Social Justice and Empowerment to address these issues, further efforts are needed to improve the coverage and quality of existing public welfare programs, ensuring they effectively target these vulnerable sections of society. These bottlenecks could be largely addressed by improving the coverage and provisions of existing public welfare programs such as Pradhan Mantri Awas Yojana (PMAY), Jal Jeevan Mission (JJM), Swachh Bharat Mission (SBM), Pradhan Mantri Sahaj Bijli Har Ghar Yojana (Saubhagya) and Pradhan Mantri Ujjwala Yojana (PMUY) or setting up of more targeted programs designed to address vulnerabilities of the population.

The Multidimensional Poverty approach, as envisaged by the UNDP, enables significant flexibility in incorporating country-specific indicators and weights. India's NITI Aayog, utilizing the Alkire-Foster methodology, innovatively added bank account (weight = 1/21) and maternal health (weight = 1/12) as additional indicators, reflecting the nation's unique developmental context (NITI Aayog, 2023). This methodological adaptation, driven by the success of large-scale policy programs such as Janani Suraksha Yojana (JSY), Pradhan Mantri Surakshit Matritva Abhiyan (PMSMA), and Pradhan Mantri Jan-Dhan Yojana (PMJDY), resulted in marginal variations in MPI calculations. While UNDP reported India's MPI for 2019–21 to be 0.069, the NITI Aayog report estimated it to be 0.066 (UNDP, 2023). Similarly, UNDP reported a poverty intensity of 42% for 2019–21, while NITI Aayog estimated it to be 44.39%. These findings indicate substantial progress in reducing multidimensional poverty in India, highlighting the effectiveness of equitable pro-poor programs and policy initiatives. The inclusion of additional indicators relevant to India's specific context provides a more comprehensive picture of poverty reduction efforts.

Targeted government interventions have been pivotal in addressing multidimensional poverty. Government interventions have been pivotal in addressing the specific deprivations tracked by the 12 MPI indicators. For Health (Nutrition, Child Mortality, Maternal Health), schemes such as Poshan Abhiyaan (Poshan 2.0), Janani Suraksha Yojana (JSY), and the Pradhan Mantri Matru Vandana Yojana (PMMVY) have directly targeted nutritional intake and maternal care coverage. In Education (Schooling and Attendance), the Samagra Shiksha Abhiyan and the Right to Education Act have been instrumental in improving enrollment and retention. The Standard of Living indicators have been addressed through targeted infrastructure missions: Pradhan Mantri Ujjwala Yojana (Cooking Fuel), Swachh Bharat Mission (Sanitation), Jal Jeevan Mission (Drinking Water), Saubhagya (Electricity), and Pradhan Mantri Awas Yojana (Housing). The Pradhan Mantri Jan Dhan Yojana (PMJDY) has specifically addressed the Bank Account indicator. Furthermore, specific interventions have been crucial for vulnerable groups. Income-enhancing schemes such as the Mahatma Gandhi National Rural

Employment Guarantee Act (MGNREGA) and the Deendayal Antyodaya Yojana–National Rural Livelihoods Mission (DAY–NRLM) have empowered rural households, particularly SCs and STs, by providing wage employment and strengthening self-help groups. For women, schemes like Stand Up India and Mahila Samman Savings Certificate promote financial independence, while the Beta Bachao Beti Padhao scheme addresses gender disparities in education, indirectly influencing future household income potential.

The Ministry of Social Justice and Empowerment has played a crucial role in improving public provisioning for marginalized communities. However, persistent multidimensional poverty remains evident among vulnerable populations. Flow characteristics suggest that continued focus on health, nutrition, and education dimensions could further reduce MPI, particularly among dependent populations and marginalized communities. For instance, the Integrated Child Development Programme played a major role in uplifting nutritional standards (Das et al., 2022). Nevertheless, continued refinement of intervention strategies remains crucial for sustained poverty reduction (Roy, 2025). Regular follow-up and evaluation of these programs are pivotal to ensure they achieve their objectives in the targeted groups. Furthermore, analysis at the administrative level is required to rigorously understand the causes of implementation gaps.

This study was subject to several methodological limitations that warrant careful consideration. The cross-sectional nature of the data inherently restricted our ability to establish any causal relationships. Methodological constraints require the omission of variables with missing values, and the reliance on self-reported data from the National Family Health Survey (NFHS) introduces potential reporting inaccuracies and bias. While our findings demonstrated broad consistency with the NITI Aayog report, slight variations arose from the nuanced methodology of indicator specification and weighting. The study was constrained to utilizing NITI Aayog's predefined indicators, predominantly focused on publicly provisioned metrics, which may not comprehensively capture the full complexity of individual poverty experiences. While this study analyzes social groups across states, it does not explicitly weight for the varying population shares of these groups within specific regions. For instance, while the share of the SC population may be proportionally higher in Punjab compared to Kerala, the economic and human development outcomes for SC households in Kerala may differ significantly due to state-specific socio-political histories. Such nuances regarding the interplay between population share and development outcomes require further granular research and were beyond the scope of this decomposition analysis. While we incorporated the two country-specific indicators (maternal health and bank accounts) to align with the NITI Aayog's national MPI, minor discrepancies in our estimates compared to the official report may persist due to slight variations in indicator operationalization within the Alkire–Foster framework. These methodological limitations highlight the need

for further research to develop more refined approaches that can comprehensively capture the multidimensional nature of poverty. Future research could benefit from incorporating more granular data collection and analysis techniques to provide a nuanced understanding of socioeconomic deprivation.

The analysis reveals a significant decline in multidimensional poverty in India, primarily driven by targeted government interventions. However, disparities persist across regions and socioeconomic groups. While the headcount ratio has shown substantial improvement, the intensity of poverty remains a concern. Several Government initiatives have made strides in addressing these issues; however, the health dimension, particularly nutrition, continues to be a significant contributor to the MPI. To further accelerate poverty reduction, a sustained focus on vulnerable groups, especially Scheduled Tribes, females, children under 14 years of age, and rural populations, is essential. Additionally, enhancing the quality and coverage of public services, promoting inclusive growth, and addressing the root causes of poverty, such as inequality and social exclusion, will be crucial for achieving equitable growth.

References

- Alkire, S., & Foster, J. (2011). Counting and multidimensional poverty measurement. *Journal of public economics*, 95(7-8), 476-487.
- Alkire, S., & Santos, M. E. (2011). Acute multidimensional poverty: A new index for developing countries.
- Alkire, S., & Seth, S. (2015). Multidimensional poverty reduction in India between 1999 and 2006: Where and how?. *World Development*, 72, 93-108.
- Alkire, S., Kanagaratnam, U., & Suppa, N. (2024). A methodological note on the global Multidimensional Poverty Index (MPI) 2024 changes over time results for 86 countries.
- Alkire, S., Oldiges, C., & Kanagaratnam, U. (2021). Examining multidimensional poverty reduction in India 2005/6-2015/16: Insights and oversights of the headcount ratio. *World Development*, 142, 105454.
- Bagli, S., & Tewari, G. (2019). Multidimensional poverty: An exploratory study in Purulia district, West Bengal. *Economic Affairs*, 64(3), 517-527. 64. 10.30954/0424-2513.2019.7.
- Bhuyan, B., Sahoo, B. K., & Suar, D. (2020). Nutritional status, poverty, and relative deprivation among socio-economic and gender groups in India: Is the growth inclusive?. *World Development Perspectives*, 18, 100180.
- Blinder, A. S. (1973). Wage discrimination: reduced form and structural estimates. *Journal of Human resources*, 436-455.
- Canudas Romo, V. (2003). Decomposition methods in demography (Unpublished internal doctoral thesis, University of Groningen). s.n.

Das, P., Ghosh, S., & Paria, B. (2022). Multidimensional poverty in India: a study on regional disparities. *GeoJournal*, 87(5), 3987–4006.

Das, P., Ghosh, S., & Paria, B. (2023). Multidimensional poverty in India: patterns of reduction across population subgroups and geographical locations during 2005–06 and 2019–21. *GeoJournal*, 88(4), 3851–3870.

Das, P., Paria, B., & Firdaush, S. (2021). Juxtaposing consumption poverty and multidimensional poverty: A study in Indian context. *Social Indicators Research*, 153(2), 469–501.

Dubey, A. (2009). Poverty and under-nutrition among scheduled tribes in India: A disaggregated analysis. IGIDR Proceedings/Project Reports Series, [from <http://www.igidr.ac.in/pdf/publication/PP-062-13.pdf>, accessed on 12-12-2013].

Government of India, Ministry of Health & Family Welfare. (2021). National Family Health Survey-5 (2019–21). <http://rchiips.org/nfhs/pdf/NFHS5/India.pdf>

IIPS. (n.d.). National Family Health Survey. International Institute for Population Sciences (IIPS). Retrieved from <http://rchiips.org/nfhsnew/nfhsuser/index.php>

Jann, B. (2008). The Blinder–Oaxaca decomposition for linear regression models. *The stata journal*, 8(4), 453–479.

Kaibarta, S., Mandal, S., Mandal, P., Bhattacharya, S., & Paul, S. (2022). Multidimensional poverty in slums: an empirical study from urban India. *GeoJournal*, 87(Suppl 4), 527–549.

Kitagawa, E. M. (1955). Components of a difference between two rates. *Journal of the american statistical association*, 50(272), 1168–1194.

Mehta, A. K., & Shah, A. (2003). Chronic poverty in India: Incidence, causes and policies. *World Development*, 31(3), 491–511.

Menon, P., Bamezai, A., Subandoro, A., Ayoya, M. A., & Aguayo, V. M. (2015). Age-appropriate infant and young child feeding practices are associated with child nutrition in India: insights from nationally representative data. *Maternal & child nutrition*, 11(1), 73–87.

NITI Aayog. (2023). National Multidimensional Poverty Index: A progress review 2023. Government of India.

Oaxaca, R. (1973). Male–female wage differentials in urban labor markets. *International economic review*, 693–709.

Oaxaca, R. L., & Sierminska, E. (2025). Oaxaca–Blinder meets Kitagawa: What is the link?. *PLoS One*, 20(5), e0321874.

Oxford Poverty and Human Development Initiative. (2018). Global Multidimensional Poverty Index 2018: The most detailed picture to date of the world’s poorest people.

Oxford Poverty and Human Development Initiative (OPHI), University of Oxford.

- Pradhan, I., Kandapan, B., & Pradhan, J. (2022). Uneven burden of multidimensional poverty in India: A caste based analysis. *Plos one*, 17(7), e0271806.
- Prasad, J. B., Pezhhan, A., & Patil, S. H. (2021). Effect of wealth, social inequality, Mother's BMI, and education level on child malnutrition in India. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews*, 15(6), 102304.
- Preston, S. H., Heuveline, P., & Guillot, M. (2001). *Demography: Measuring and modeling population processes*. Blackwell Publishers Ltd.
- Raghunathan, K., Headey, D., & Herforth, A. (2021). Affordability of nutritious diets in rural India. *Food Policy*, 99, 101982.
- Rahimi, E., & Hashemi Nazari, S. S. (2021). A detailed explanation and graphical representation of the Blinder–Oaxaca decomposition method with its application in health inequalities. *Emerging Themes in Epidemiology*, 18(1), 12
- Ramachandran, P. (2010). Nutrition and child survival in India. *The Indian Journal of Pediatrics*, 77, 301–305.
- Rogan, M. (2016). Gender and multidimensional poverty in South Africa: Applying the global multidimensional poverty index (MPI). *Social Indicators Research*, 126, 987–1006.
- Roy, A. (2025). Multidimensional Poverty and Quality of Governance in Indian States. *The Indian Economic Journal*, 73(3), 548–566.
- Roy, P., Ray, S., & Haldar, S. K. (2019). Socio-economic determinants of multidimensional poverty in rural West Bengal: A household level analysis. *Journal of Quantitative Economics*, 17(3), 603–622.
- Rupavath, R. (2023). Poverty and education: Attainments and challenges for tribal communities. *Contemporary Voice of Dalit*, 15(2), 230–247.
- Ryckman, T., Beal, T., Nordhagen, S., Murira, Z., & Torlesse, H. (2021). Affordability of nutritious foods for complementary feeding in South Asia. *Nutrition reviews*, 79(Supplement_1), 52–68.
- Sahoo, P., Mondal, S., & Kumar, V. (2023). Multidimensional deprivations among social groups in rural India: A state level analysis. *GeoJournal*, 88(6), 6137–6159.
- Sen, A. K., & Anand, S. (1997). Concept of human development and poverty: A multidimensional perspective. In *Poverty and human development papers 1997* (pp. 1–20). United Nations Development Programme.
- Singh, A. (2019). Social Justice and Empowerment for Marginalized Groups: Realities and Responses. *Global Initiatives for Sustainable Development: Issues and Strategies*, 184.
- Tripathi, S., & Yenneti, K. (2020). Measurement of multidimensional poverty in India: A State-level analysis. *Indian Journal of Human Development*, 14(2), 257–274.

UNDP. (2023). 2023 Global Multidimensional Poverty Index (MPI): Unstacking global poverty: Data for high impact action. United Nations Development Programme.

United Nations Children's Fund. (2024). Child food poverty: Nutrition deprivation in early childhood (Child nutrition report, June 2024). UNICEF.

United Nations Development Programme. (1997). Human development report 1997. <http://hdr.undp.org/en/content/human-development-report-1997>

United Nations. (2024). Global indicator framework: SDG indicators. <https://unstats.un.org/sdgs/indicators/Global-Indicator-Framework-after-2024-refinement-English.pdf>

Welthungerhilfe (WHH), Concern Worldwide, & Institute for International Law of Peace and Armed Conflict (IFHV). (2024). 2024 Global Hunger Index: How gender justice can advance climate resilience and zero hunger. WHH; Concern Worldwide; IFHV.

Prevalence and Correlates of Overweight and Obesity in Adults and Older Adults in India; Population-Level Estimates Based on Nationally Representative Surveys (2015–21)

Debayanti Bhowmick[#], Abhishek Kumar^{*}, Ajay Kumar Verma[^]

In the current epidemiological transition, a rapid increase in the prevalence of overweight is observed in low-middle-income countries. There is a prominent shift towards a sedentary lifestyle and unhealthy food consumption which has triggered the risk of NCDs. The situation is similar in India. A holistic understanding of obesity epidemiology is the need of the hour with a target-oriented intervention approach. Though multiple studies have been a part of the research pool, insights are limited in the case of the whole prevalence as per the National estimate. This study attempts to understand the epidemiology of overweight/obesity in India concerning the entire population. The study is unique in the sense that it utilizes National-level estimates to understand the epidemiology of obesity concerning the entire population. It aims to explore the state-level prevalence of overweight/obesity and understand the implications as per socio-economic indicators and age-sex groups. The study takes up cross-sectional analysis, using two National Representative data sets. It identifies key anthropometric parameters and socio-economic correlates to run descriptive analysis and logistic regression. The descriptive analysis provided us with the age-sex prevalence of obesity, while logistic regression was used to establish the SES gradient by age. Lastly, to measure the inequality of obesity among males and females by different age groups, concentration indices is used. The findings reveal a higher prevalence of overweight/obesity among females than males. The prevalence among females is 19.2 percent, whereas it is 16.8 percent in the case of males. The highest prevalence is observed in NCT Delhi (33.9%) and the lowest in Meghalaya (10%). With respect to socio-economic determinants, the prevalence is noticeably higher among those residing in urban areas, belonging to the richest wealth index, identifying in the 'others' social groups and who have 10 or more years of education across the age groups. The paper presents all-inclusive results of obesity/overweight epidemiology in India, giving a holistic understanding of obesity implications with regard to age and SES-wise determinants. Thus, providing probable pathways for intervention.

Keywords: Obesity, Overweight, India, Nutrition, Body Mass Index

[#] Corresponding Author: Debayanti Bhowmick, University of New South Wales, Sydney, Australia
Email: d.bhowmick@unsw.edu.au

^{*} FLAME University, Pune, India

[^] Banaras Hindu University, Varanasi, India

The rapid increase in the prevalence of overweight/obesity in low- and middle-income countries (LMICs) is a more relatively recent trend globally (McLachlan & NCD-RisC, 2016). An increase has been noted in both rural and urban areas, with the rural areas displaying a recent pattern of intensified escalation (NCD-RisC, 2019). The rise in overweight and obesity rates across age groups has been relatively higher in LMICs and plateauing in high-income countries (HICs). As of 2016, over two billion (70%) globally overweight or obese individuals, reside in LMICs (Abay et al., 2022). While the major direct cause is the rapid increase in consumption of ultra-processed food along with other junk foods high in added saturated fats, sodium and sugar, underlying factors include rapid social and economic changes such as urbanisation, globalization, and economic growth accompanied by increased income per capita (Wen et al., 2009). Increased sedentary living occurred earlier and now it is the diet that directly impacts these changes (Popkin & Ng, 2022). Certain subpopulations also experience the dual challenge of malnutrition, resulting in both hunger-induced undernutrition and stunting. The rising consumption of ultra-processed food and beverages, and increased prevalence of overweight and obesity have triggered the risks of multiple Non-Communicable Diseases (NCDs) such as cardiovascular diseases (CVDs), diabetes, hypertension and 13 of the 16 major cancers (WCRF & AICR, 2018). The increased mortality from CVD and cancer is also very concerning (Popkin, 2003). Over the last few decades, the prevalence of NCDs has sharply increased in many LMICs, while it is somewhat declining in the HICs (Abdelaal et al., 2017).

India is going through a similar nutrition transition (Popkin & Ng, 2022). Amidst rapid economic growth and income development, the prevalence of overweight/obesity and associated NCDs has been consistently increasing across all age groups and regions in India (Miranda et al., 2019). Moreover, the figures are expected to triple among Indian adults (20–69 years) between 2010 and 2040 with higher risk among the rural residents and older Indians, if not intervened timely (Ranjani et al., 2016). This is mainly linked with the rapid increase in consumption of ultra-processed food by the Indian population, particularly those under 40 years of age.

The cross-sectional analysis presented in this study has a two-fold advantage. Firstly, it provides evidence on the age group where the rise of obesity prevalence is triggered. Secondly, when compared with lagged estimates across the age-sex group, it can provide an understanding of metabolic problems accumulating over time. More importantly, the paper is an insight into the reliability of the outcomes as obtained from using 2 different data sets positioned in different time frames. Though the issue being addressed here is very pertinent to public health, the current data repository on obesity may not potentially produce a reliable glimpse of the actual epidemiological status. It is established that the pattern of BMI changes in accordance with age (Luhar et al., 2020). Thus, for ideal quantification, taking up a life course pattern is critical to understanding the magnitude of the burden.

Currently, the literature on prevalence and correlates has mostly focused on socioeconomic status (SES) and its association with overweight and obesity. Studies have reported a positive association between SES indicators and obesity across age groups (Neuman et al., 2013). However, evidence suggests that the association vary by economic development, region, and gender within countries (Monteiro et al., 2004). There is a complex linkage between SES levels and obesity. While decades ago, SES in LMICs was linked positively with obesity, more recently the shifts have been complex and many countries such as China, Indonesia, and others have higher levels of obesity among the poor than the higher SES groups (Popkin et al., 2020).

In LMICs, a positive association is likely across age groups and sex (adult men and women and children), while in developed countries, a negative association is more common among women (inconsistent in men and children) (Pavela et al., 2016). A recent review by Templin et al., (2019), reveals that as national economies increase to higher levels, the burden of overweight/obesity shifts over time to poorer sections of the population within these countries. The impact factor shows a considerable increase in prevalence amongst the poorer sections of the population whilst, for the wealthiest sub-populations, the prevalence rate remains more or less unchanged (Cohen et al., 2013).

The present paper answers the following research questions specifically for adults and older adults using India's two nationally representative surveys: (a) What is the prevalence of overweight/obesity in India? (b) Are wealthy/educated more likely to be overweight/obese in India? (c) Whether thinning set in earlier for men than women? (d) Whether the current data set is reliable for quantifying the magnitude of obesity in India?

The question aims to understand the epidemiology of overweight/obesity in India concerning the entire population. It takes into consideration the respective age-sex groups, their experiences and socio-economic status. To round up the analysis and its implications we also explore the state-level associations of overweight/obesity prevalence.

Methodology

• Study Design

The study is a cross-sectional analytical study using two National Representative data sets. It identifies anthropometric parameters and socio-economic correlates to run descriptive analysis and logistic regression. Additionally, an "Obesity Tree/Pyramid" depicting age-sex-wise prevalence was also plotted.

• Variables

The main variables used are the anthropometric parameters to estimate the prevalence of overweight and obesity. WHO adult BMI cut-offs, BMI for age Z score (BAZ), and weight-for-height Z-score (WHZ) as per WHO standard are used for adults respectively. Body mass index (BMI) is defined as a person's weight in kilograms divided by the square of the height in meters (kg/m²) that is commonly used to classify overweight and obesity in adults (Pavla et al., 2016)..

For adults (all ages), WHO defines overweight as BMI ≥ 25 and obesity as BMI ≥ 30 (Templin et al., 2019).

Alternate measures to identify abdominal obesity in adults i.e., waist-to-hip ratio were also used. In accordance with the WHO, a waist-to-hip ratio above 90 cm in men and >85 cm in women is classified as abdominal obesity.

Other background variables used in the analysis consisted of a few correlates including socioeconomic factors including i) place of residence (urban/rural); ii) wealth quintiles as poorest, poor, middle, rich and richest; iii) social groups as SC, ST, OBC and 'Others' and iv) education in years as <5 years, 5–9 years and 10 years and more.

• Data Sources

The study is based on nationally representative surveys namely National Family Health Survey (NFHS), 2019–21, and Longitudinal Ageing Study in India (LASI) Wave 1, 2020. A brief overview of the surveys is given below.

- National Family Health Survey, 2019–21

The National Family Health Survey, 2019–21 survey was conducted by International Institute for Population Sciences (IIPS), Mumbai across all India's States and UTs using a two-stage, stratified cluster sample, wherein 724,115 women (15–49 years) and 101,839 men (15–54 years) were selected from a random sample of 636,699 households. The survey yielded an analytical sample of 423,843 adult women (20–49 years) and 65,196 men (20–49 years) after adjusting missing data which were used in the present analysis. For more information on the sampling procedure of the NFHS–2019–2021 survey, visit the link http://rchiips.org/nfhs/NFHS-5Reports/NFHS-5_INDIA_REPORT.pdf

- Longitudinal Ageing Study in India (LASI) Wave 1, 2017–19

Longitudinal Ageing Study in India (LASI) 2020 (Wave-I) is a nationally demonstrative survey of the health, economic, and social determinants and consequences of older adults (45 and above years) and population ageing in India. The LASI Wave 1 was conducted across India's States and UTs (excluding Sikkim) by International Institute for Population Sciences (IIPS), Mumbai using a stratified,

multistage stratified area probability cluster sampling design wherein 72,250 older adults and elderly men and women aged 45 and above were selected.

The present study used 45105 observations after adjusting missing data of the older population who were 50 and above years. For more information on the sampling design and method of the LASI-2020 (Wave-I) survey, visit the link <https://www.iipsindia.ac.in/content/lasi-publications>, <https://www.iipsindia.ac.in/content/lasi-publications>.

• Statistical methods

Descriptive statistics were conducted to estimate prevalence by correlates. To estimate the state-level prevalence of obesity, NFHS-4 data were used for 15-49 years, and LASI datasets for 50-59 years, 60-69 years and 70 years and above. The weighted prevalence population data for 2021 was obtained from the Population Projection Report 2011-2036. The share for each age group was calculated in the total population which served as weights to calculate the weighted prevalence. This was done at both the national and the State levels.

Logistic regression analyses were carried out; the odds ratio for correlates to establish the SES gradient by age. Interactions of wealth quintile/education to examine the SES gradient by age have also been made. To measure the inequality of obesity among males and females by different age groups we have used the concentration indices.

Results

Among the male population, the prevalence of obesity increases with age up to 40-49 years and starts to decline thereafter. Whereas, among females, the percentage of obesity increases up to the age of 50-54 years and declines after that. The overall prevalence of overweight and obesity is 17 percent among 20-29 years, 31.7 percent among 30-39 years, 36.4 percent among 40-49 years, 32 percent among 50-59 years, 25.3 percent among 60-69 years and 16.9 percent among ≥ 70 years (Table 1). As per the SES determinants, the prevalence is comparatively higher among those belonging to urban areas, the richest community, 'others' social groups and higher education experience. (Table 1). Hereafter we use the term obesity but it combines both overweight and obesity. We combined the figures as in India the risk of diabetes and many other NCDs rises at a BMI of 22 (Wen et al., 2009).

The prevalence of obesity is highest in Delhi (33.9 %) with females accounting for 37.3 percent and males accounting for 30.9 percent. The prevalence is lowest in Meghalaya (10 %) with females having a 9.9 percent prevalence rate and males recording 10.5 percent (Table 2).

Table 1: Age-group-wise prevalence of overweight and obesity by SES for total population

Background Variables	20-29 Years	30-39 Years	40-49 Years	50-59 Years	60-69 Years	>=70+ Years
Place of residence	%	%	%	%	%	%
Urban	22.8	41.9	48.3	52.4	43.3	32.6
Rural	14.3	26.6	30.3	22.8	18.2	11.1
Wealth Index						
Poorest	7.3	13.7	16.0	23.1	16.3	10.6
Poor	12.1	22.8	24.8	27.2	21	13.1
Middle	17.0	31.7	35.1	29.4	25	17
Rich	21.2	40.4	44.4	35.3	31.9	19.4
Richest	26.0	47.7	55.9	47.4	35.4	27.6
Social Groups						
SC	15.7	29.4	32.6	22.8	19.2	9.1
ST	9.1	17.5	20.1	15.4	8.8	5.5
OBC	17.4	32.8	37.7	36	26.2	19
Others	20.4	37.1	42.3	38.7	34.2	22
Education in years						
No schooling	10.3	20.6	26.9	23.4	17.9	11.3
<5 years	12.3	24.9	33.3	24.2	24.1	19
5-9 years	16.0	32.6	41.1	37.6	32.8	20.2
10+ years	19.1	40.9	49.0	51.3	42.2	39.5
Total	17.0	31.7	36.4	32	25.3	16.9

Note- N is the weighted total sample.

The values of concentration indices are positive across all age groups indicating that overweight/obesity is concentrated among the wealthy population as compared to the poor (Table 3).

It is observed that the odds of obesity were greater for urban dwellers as compared to their rural counterparts. For urban adults aged 20-29 and 30-39 years, odds are 1.18 (95% CI: 1.15-1.21) and 1.14 (95% CI: 1.11,1.16) times the odds of obesity. The figure is similar for adults aged 40-49 years i.e., 1.20 times. The odds go comparatively higher for older adults and the elderly. Within the age group 50-59, 60-69 and 70+, there are 2.61 (95%, CI: 2.43, 2.80), 2.64 (95% CI: 2.45, 2.86) and 2.86 (95% CI: 2.56, 3.19) times the odds of obesity, respectively (Table 4). At the same time, it is important to note the rapid increases in rural obesity over the past decade (NCD- RisC, 2019).

As per the wealth index, the odds of obesity are higher for the wealthier population. This section states the odds as compared between the poorest and the richest. The richest adults belonging to age groups 20-29, 30-39 and 40-49 years had 3.93

Table 2: State standard estimates of obesity and overweight

State/UT	Male %	Female %	Total %
Andhra Pradesh	25.7	31.9	28.8
Assam	13.9	13.4	13.7
Bihar	11.5	11.8	11.6
Chhattisgarh	12.7	11.4	12.1
Gujarat	16.8	21.9	19.2
Haryana	21.5	25.7	23.5
Himachal Pradesh	23.9	29.1	26.5
Jammu & Kashmir	24.8	27.9	26.2
Jharkhand	11.1	9.6	10.4
Karnataka	26.1	30.1	28.1
Kerala	28.0	34.7	31.4
Madhya Pradesh	12.4	13.4	12.8
Maharashtra	20.5	22.7	21.5
NCT Of Delhi	30.9	37.3	33.9
Odisha	18.0	17.7	17.2
Punjab	27.9	37.0	32.2
Rajasthan	12.2	12.5	12.4
Tamil Nadu	27.6	33.9	30.7
Telangana	23.4	24.9	24.1
Uttar Pradesh	13.8	16.3	15.0
Uttarakhand	21.9	24.3	23.1
West Bengal	15.6	18.8	17.1
Arunachal Pradesh	20.9	19.4	20
Goa	26.8	32.9	30
Manipur	23.3	26.7	25
Meghalaya	10.5	9.9	10
Mizoram	22.6	17.9	20
Nagaland	18.9	14.5	17
Sikkim	NA	NA	NA
Tripura	18.5	17.1	18

(95% CI: 3.70, 4.07), 4.41 (95% CI: 4.23, 4.58) and 4.95 (95% CI: 4.84, 5.25) times the odds of obesity, respectively. In the same wealth index, older adults and the elderly aged 50–59, 60–69 and 70+ years accounted for comparatively lesser odds of obesity i.e., 2.38 (95% CI: 2.14, 2.66), 2.36 (95% CI: 2.09, 2.66) and 2.44 (95% CI: 2.04, 2.91), respectively.

Concerning social identity, it is observed that the likelihood of obesity in the category of ‘others’ is higher than those belonging to the Scheduled Caste category.

Table 3: Concentration index values for the total population

Age group	Index value	Std. error	p-value	No. of obs.
20 to 29 years	0.23	0.00	0.00	232154
30 to 39 years	0.23	0.00	0.00	212355
40 to 49 years	0.23	0.00	0.00	180963
50 to 59 years	0.14	0.01	0.00	18225
60 to 69 years	0.15	0.01	0.00	16529
70+ years	0.18	0.01	0.00	10351

Table 4: Logistic regression on overweight and obesity by SES for total population

Background Variables	20-29 Years	30-39 Years	40-49 Years	50-59 Years	60-69 Years	>=70+ Years
Place of residence						
OR [95%, C.I.]						
Rural	1	1	1	1	1	1
Urban	1.18***	26.6	30.3	22.8	18.2	11.1
Wealth Index						
Poorest	1	1	1	1	1	1
Poor	"1.69*** [1.62, 1.77]"	"1.77*** [1.71, 1.84]"	"1.66*** [1.60, 1.72]"	1.19** [1.07, 1.33]	1.26*** [1.11, 1.42]	"1.29** [1.08, 1.54]"
Middle	"2.42*** [2.31, 2.53]"	"2.59*** [2.50, 2.68]"	"2.50*** [2.41, 2.59]"	1.33*** [1.19, 1.49]	1.51*** [1.34, 1.71]	1.52*** [1.27, 1.81]
Rich	"3.10*** [2.95, 3.25]"	"3.43*** [3.30, 3.56]"	"3.43*** [3.29, 3.56]"	1.72*** [1.55, 1.92]	1.96*** [1.74, 2.21]	2.08*** [1.75, 2.48]
Richest	3.93***[3.74, 4.14]	4.41***[4.23, 4.60]	4.95***[4.74, 5.17]	2.38*** [2.14, 2.66]	2.36*** [2.09, 2.66]	2.44*** [2.04, 2.91]
Social Groups						
SC	1	1	1	1	1	1
ST	"0.82*** [0.79, 0.85]"	"0.75*** [0.73, 0.78]"	"0.82*** [0.79, 0.85]"	0.72*** [0.64, 0.81]	0.77*** [0.67, 0.88]	0.81 [0.66, 1.00]
OBC	"1.01 [0.98, 1.04]"	"0.95*** [0.93, 0.98]"	"0.99 [0.96, 1.02]"	1.18*** [1.07, 1.30]	1.25*** [1.12, 1.39]	"1.21* [1.02, 1.43]"
Others	"1.17*** [1.13, 1.21]"	"1.09*** [1.06, 1.12]"	"1.11*** [1.08, 1.15]"	1.49*** [1.34, 1.65]	1.47*** [1.30, 1.65]	1.49*** [1.25, 1.78]
Education						
No schooling	1	1	1	1	1	1
<5 years	"1.08 [1.00, 1.17]"	"1.12*** [1.07, 1.18]"	"1.17*** [1.13, 1.22]"	"1.00 [0.90, 1.13]"	1.20** [1.07, 1.36]	"1.20* [1.03, 1.41]"
5-9 years	"1.18*** [1.13, 1.24]"	"1.30*** [1.26, 1.33]"	"1.28*** [1.25, 1.32]"	1.43*** [1.32, 1.56]	1.45*** [1.32, 1.59]	"1.22** [1.05, 1.40]"
10 years & more	"1.10*** [1.05, 1.15]"	"1.33*** [1.29, 1.37]"	"1.34*** [1.30, 1.38]"	1.59*** [1.44, 1.74]	1.69*** [1.53, 1.88]	1.53*** [1.31, 1.78]

Note- P <0.01***, P <0.05** and *<0.10

Table 5: Logistic regression of Interactions between years of schooling and wealth index for total population on obesity

Schooling years and wealth	20–29 years OR [95%, C.I.]	30–39 years OR [95%, C.I.]	40–49 years OR [95%, C.I.]	50–59 years OR [95%, C.I.]	60–69 years OR [95%, C.I.]	70+ years OR [95%, C.I.]
No school & poorest	1	1	1	1	1	1
No school & poorer	1.77***[1.59, 1.96]	1.96***[1.86, 2.08]	1.71***[1.63, 1.80]	1.15 [0.98, 1.35]	1.32** [1.11, 1.56]	1.39** [1.10, 1.76]
No school & middle	2.64***[2.35, 2.96]	2.92***[2.75, 3.10]	2.68***[2.56, 2.82]	1.41*** [1.21, 1.66]	1.74*** [1.47, 2.06]	1.73*** [1.37, 2.17]
No school & richer	3.96***[3.47, 4.53]	4.45***[4.15, 4.77]	3.88***[3.68, 4.09]	1.81*** [1.55, 2.12]	2.15*** [1.81, 2.54]	2.12*** [1.69, 2.66]
No school & richest	4.87***[4.02, 5.90]	5.98***[5.41, 6.61]	6.08***[5.69, 6.51]	2.38*** [2.02, 2.80]	2.62*** [2.20, 3.12]	2.74*** [2.16, 3.47]
<5 years & poorest	1.06[0.91, 1.23]	1.20***[1.09, 1.32]	1.07[0.97, 1.19]	1.17 [0.90, 1.51]	1.61*** [1.22, 2.11]	1.71** [1.18, 2.46]
<5 years & poorer	1.95***[1.69, 2.25]	2.05***[1.88, 2.24]	2.07***[1.91, 2.24]	1.28* [1.01, 1.64]	1.95*** [1.52, 2.50]	1.76** [1.23, 2.52]
<5 years & middle	2.86***[2.43, 3.37]	3.42***[3.13, 3.74]	3.16***[2.93, 3.42]	1.56*** [1.23, 1.98]	2.25*** [1.76, 2.89]	1.96*** [1.37, 2.80]
<5 years & richer	4.03***[3.31, 4.90]	4.68***[4.20, 5.23]	4.68***[4.29, 5.10]	1.82*** [1.44, 2.29]	3.00*** [2.33, 3.87]	3.13*** [2.27, 4.31]
<5 years & richest	6.56***[4.86, 8.86]	6.39***[5.32, 7.68]	6.99***[6.12, 7.98]	3.02*** [2.38, 3.83]	3.15*** [2.43, 4.09]	4.48*** [3.26, 6.14]
5 9years & poorest	1.16**[1.06, 1.28]	1.41***[1.32, 1.50]	1.34***[1.24, 1.45]	2.02*** [1.68, 2.44]	2.42*** [1.95, 3.01]	2.29*** [1.61, 3.26]
5 9years & poorer	2.08***[1.91, 2.26]	2.45***[2.32, 2.58]	2.16***[2.05, 2.29]	2.41*** [2.03, 2.86]	2.87*** [2.35, 3.50]	2.56*** [1.88, 3.48]
5 9years & middle	3.27***[3.00, 3.55]	3.84***[3.65, 4.04]	3.42***[3.25, 3.60]	2.66*** [2.24, 3.16]	2.88*** [2.37, 3.50]	2.35*** [1.73, 3.20]
5 9years & richer	4.29***[3.93, 4.67]	5.31***[5.04, 5.60]	4.97***[4.73, 5.23]	3.36*** [2.85, 3.96]	3.70*** [3.05, 4.48]	3.62*** [2.73, 4.82]
5 9years & richest	5.57***[5.05, 6.15]	7.33***[6.91, 7.79]	7.91***[7.48, 8.35]	3.92*** [3.32, 4.62]	4.32*** [3.57, 5.24]	3.60*** [2.67, 4.85]
10+more & poorest	1.21***[1.09, 1.35]	1.73***[1.56, 1.93]	1.49***[1.28, 1.73]	3.39*** [2.66, 4.33]	3.72*** [2.81, 4.91]	2.97*** [1.87, 4.73]
10+more & poorer	2.02***[1.85, 2.19]	2.80***[2.62, 2.99]	2.53***[2.33, 2.75]	3.42*** [2.78, 4.21]	4.27*** [3.37, 5.42]	3.75*** [2.58, 5.44]
10+more # middle	2.86***[2.64, 3.10]	3.87***[3.67, 4.09]	3.79***[3.56, 4.04]	3.23*** [2.67, 3.92]	4.49*** [3.64, 5.54]	4.56*** [3.30, 6.31]
10+more& richer	3.89***[3.60, 4.20]	5.25***[5.00, 5.52]	5.24***[4.97, 5.52]	4.34*** [3.66, 5.15]	5.35*** [4.40, 6.52]	6.17*** [4.64, 8.22]
10+more & richest	5.36***[4.97, 5.79]	7.32***[6.98, 7.67]	8.01***[7.65, 8.39]	5.79*** [4.96, 6.76]	6.71*** [5.63, 8.01]	6.24*** [4.85, 8.01]

Note– P <0.01***, P <0.05** and * <0.10

Adults in the 'others' category, aged 20–29, 30–39 and 40–49 years have 1.17 (95% CI: 1.12, 1.20), 1.09 (95% CI: 1.06, 1.12) and 1.11 (95% CI: 1.08, 1.15) times higher odds of obesity, respectively. Lastly, older adults and elderly aged 50–59 (95% CI: 1.34, 1.65) and 70+ years (95% CI: 1.25, 1.78) have 1.47 times and 1.49 times the odds of obesity, respectively.

Education being the final determinant, the propensity of obesity is higher among the population having ten or more years of education. Obesity holds 1.10 times higher odds among adults aged 20–29 years (95% CI: 1.08, 1.18), having ten years and more education. The odds are 1.33 and 1.34 for age groups 30–39 and 40–49 years, respectively, within the same education parameter. Among the older adults and the elderly, the population aged 50–59, 60–69 and 70+ years have 1.59 (95% CI: 1.44, 1.74), 1.69 (95% CI: 1.53, 1.88) and 1.53 (95% CI: 1.31, 1.78) times the odds of obesity.

Table 5 shows the logistic regression between the level of education (in years) and the wealth index on obesity for the total population. Interaction analysis reports that those with an education of ten or more years and belonging to the richest section were significantly more likely to be obese as compared to those who had no education and belonged to the poorest wealth index.

Discussion

According to the available data, this study is a one-of-a-kind insight into the epidemiology of obesity/overweight in India. It uses two nationally representative surveys. This paper also provides India-based evidence on the association of overweight/obesity with SES indicators (income and education) by sex.

• Key Findings

The analysis reveals that the national prevalence of obesity across age groups is 19.2 percent in females which is sufficiently higher than in males who record a figure of 16.8 percent. The highest prevalence is observed in NCT Delhi and the lowest in Meghalaya. The prevalence is noticeably higher among those residing in urban areas, belonging to the richest wealth index, identifying in the 'others' social groups and who have 10 or more years of education across the age groups.

Thus, it is evident from the findings that the prevalence of overweight/obesity among females is higher than in males. In adults and adolescents in developing countries, a higher prevalence of obesity was seen among females, while in developed countries, the case was essentially the opposite (Mistry & Puthussery, 2015; Garrido-Miguel et al., 2019). This study estimates national overweight/obesity prevalence across all age groups. There are twofold advantages to estimating the national level prevalence i.e.; (a) It reveals a particular age group which requires priority focus

by the national programs and policies and (b) It provides an understanding of the metabolic factors accumulating over time when compared with lagged estimates across the age-sex groups.

A consistent gender disadvantage in obesity in adulthood was observed in our study. This is in line with the findings from the studies from other LMICs, where they found a much higher prevalence of obesity in women aged 20 years and above than in men (Skinner et al., 2018). The most common explanations regarding gender disparities in overweight and obesity were physical activity, cultural values, biological factors (e.g., menopause), and urbanization (Mistry & Puthussery, 2015). In contrast, Ameye & Swinnen (2019) suggests an absence of a significant gender difference in average obesity prevalence in high-income level countries. Additionally, there's a reversal of the obesity gap in high-income level countries, with males becoming the more obese gender in Japan, Scandinavian countries, and in Northern European countries like Belgium, France, Switzerland and Germany (Kanter & Caballero, 2012).

A positive SES gradient was observed for females where the prevalence of obesity increased with increasing education level and wealth quintile, which is also in line with earlier evidence from India (Bhurtyal & Adhikari, 2022). Indian studies have linked this to cultural norms that may favour fat body shapes, higher consumption of energy-dense diets, and traditional narratives which bars high-SES women in India to engage in physical activities and healthy dietary practices, despite more knowledge, awareness and resources (Kanter & Caballero, 2012). Our finding slightly deviates from the findings in developing countries. It has been observed that the women with higher SES throughout their life, have lower BMI, and the findings among men were less consistent (Ameye & Swinnen, 2019). These could be attributed to having weight-related standards among wealthy females resulting in regular exercise and healthy diets, which are of course easier to maintain with higher income.

A meta-analysis of prospective cohort studies found that individuals with obesity were associated with a 7- and 3-times higher risk of diabetes and being overweight respectively, as compared to normal-weight individuals (Abdullah et al., 2010). Our analysis found positive associations among overweight/obesity, NCD burden and out-of-pocket health expenditure. Therefore, addressing the growing overweight and obesity prevalence is of great urgency. Preventive measures such as screening for NCDs like diabetes and hypertension among overweight/obese individuals especially among high-risk individuals should be integrated into the health system. The existing government initiatives such as the National Multisectoral Action Plan for Prevention and Control of NCDs (2017-22), which partly aims to reduce out-of-pocket expenditure on NCD healthcare should be implemented across India (Luhar et al., 2020)..

This section not only elaborates on the limits but also puts forth essential suggestions significant for obesity-related research. There are multiple issues which need to be kept in mind while considering the findings of this study. Firstly, the survey took into consideration a basic method of measurement i.e., BMI (derived from weight and height) to measure the prevalence of overweight and obesity. BMI is a measurement of only relative body weight, so it does not distinguish between body fat and lean mass. There are many other better alternate anthropometric measurements such as waist circumference and waist-to-hip ratio that are strongly associated with CVD risk factors (Goh et al., 2014) along with skinfold thickness which predicts body fat.

Secondly, we have used WHO's standard global adult cut-offs i.e., ≥ 25 kg/m² for overweight and BMI ≥ 30 for obesity. However, some studies have suggested that the proposed BMI standards aren't appropriate and the recommendation is to use lower BMI cut-offs for overweight/obese (≥ 23) and obese (≥ 25) for South Asians (Misra et al., 2009). This is because South Asians are at risk of developing obesity-related co-morbidities at lower levels of body mass index (BMI) and waist circumference (WC). Yet, we have chosen to use global cut-offs in order to ease the direct comparison with other studies.

Thirdly, the rising threat of obesity is alarming and a comprehensive understanding of it is thus important and relevant. The situation is equally precarious in India, and research minds have expressed the need for integrated data repositories for fruitful outcomes and forecastable status quo (world obesity report- India). The paucity of data at the population level makes estimation even more challenging. In spite of the impacts being significant population-wise, survey methods and data repositories have confined themselves to targeted age groups be it the children, elderly or reproductive women. Like poverty, obesity is associated with multiple causalities and consequences including social and economic points of view (Egger et al., 2012). However, though the former approaches the issue through National-level holistic estimates, the latter is mainly assessed along certain target groups. The similarity of the context of both issues suggests the adoption of alike approaches- in this case, using population-level estimates. This study intends to do the same by using the available datasets across each age group. However, the datasets being disintegrated are difficult to standardise. The standard deviation calculated for each State (Table S7) shows a wide range of dispersion ranging between 0.39 (Jharkhand) to 10.78 (Nagaland) among males and 0.19 (Madhya Pradesh) to 6.87 (Nagaland) among females. The high variability of the values questions the consistency of the outcomes, especially in order to generate meaningful population-level insights. The pooled data also had minor year variations.

The appropriateness of BMI as the standard measure for obesity across age groups has been a persistent topic of deliberation. Multiple studies note how physiological status among humans may vary with ageing, making obesity measurement

challenging (Goran, 1998). It is observed that ageing is associated with decreasing body weight and height. There is also the redistribution of adipose tissues along with a decrease in muscle mass (Batsis et al., 2016). These added complexities make it difficult to accurately diagnose obesity among the elderly, especially through the BMI index. In the study done by Batsis et al. (2016), the authors strategically emphasised the diagnostic inaccuracy of BMI application on elderly populations. Their findings show that BMI may fail to interpret adiposity, which is an important determinant among elderly populations. In a systematic and meta-analytical study carried out by Correa et al. (2016), the findings reveal that out of the 13 chosen articles, 5 manuscripts present evidence of the waist-to-height ratio (WHtR) being the “best anthropometric index” when applied alone.

The study also drew an association between obesity assessed via WHtR and the capacity to predict NCD risk factors, proving its efficiency greater than BMI. Similar criticism was obtained in the study conducted by Perissinotto et al. (2002). The cross-sectional analysis revealed the drawbacks associated with the standardization of homogenous anthropometric measurements to represent every age group. The paper investigates the cross-sectional sample of an elderly population and results show marked redistribution of body fat among the elderly. Since BMI does not quite reflect this redistribution, the homogenous BMI standards misinterpret obesity among elderly subjects. Ageing is usually coupled with an increase in fat mass and a decrease in fat-free mass. Thus, the threshold values in the case of the elderly population need careful re-considerations.

Despite the ongoing criticism, BMI continues to be the most frequently used index. It is not only an easy tool but has also predicted adverse outcomes in global scenarios. Though multiple studies have recommended alternate measurement indices, BMI along with waist circumference (WC) continue to receive special attention for analytical purposes. In fact, the systematic review by Correa et al (2016) also included two such studies which recognized WHtR, WC and BMI as having the same performance level.

Fourth is a limitation associated with interpreting cross-sectional association as one cannot draw any conclusions regarding causality out of the cross-sectional association.

Conclusion

In conclusion, the findings from our analysis show a higher prevalence of obesity in women compared to men, especially among those with higher socioeconomic status. The prevalence is also more pronounced among women and adults. This indicates that India is experiencing a rapid increase in overweight/obesity prevalence.

In addition, the presence of gender disadvantage (high prevalence in adult women) and positive SES gradient for females indicate where interventions should be targeted. Rapid urbanisation and economic growth, leading to shifting towards unhealthy food consumption, fatty diets, free sugars and/or salt and a sedentary lifestyle are the major causes of overweight and obesity globally. Therefore, implementing comprehensive food policies as recommended by the World Health Organization such as food and beverage taxes, marketing restrictions on unhealthy foods and beverages, and mandatory Front-of-Pack Labelling (FOPL) system with warning labels (which has worked wonders in Chile and Mexico, and among many other countries), should be implemented in India to control the obesogenic environment and promotion of healthy dietary habits.

All age groups in India are increasingly consuming excessive ultra-processed food which foretells a future increase in overweight/obesity and all nutrition-related NCDs. The strengthening, scaling up and redesigning of existing undernutrition-related interventions, programmes and policies can be done to tackle the growing burden of overweight and obesity. It is important to realize the increasing intake of ultra-processed food among infants and toddlers. A cost-effective approach known as double duty actions may be applied. Most importantly, we must find a way to reduce intake of ultra-processed food consumed by preschoolers (Pries et al., 2019). Thus, policies and programs in low and middle-income countries must prioritize safeguarding child nutrition through increased micronutrient intake and capping the heightened distribution of ultra-processed food and beverages.

References

- Abay, K. A., Ibrahim, H., & Breisinger, C. (2022). Food policies and obesity in low- and middle-income countries. *World Development*, 151, 105775.
- Abdelaal, M., le Roux, C. W., & Docherty, N. G. (2017). Morbidity and mortality associated with obesity. *Annals of translational medicine*, 5(7), 161.
- Abdullah, A., Peeters, A., de Courten, M., & Stoelwinder, J. (2010). The magnitude of association between overweight and obesity and the risk of diabetes: a meta-analysis of prospective cohort studies. *Diabetes research and clinical practice*, 89(3), 309–319.
- Ameye, H., & Swinnen, J. (2019). Obesity, income and gender: the changing global relationship. *Global Food Security*, 23, 267–281.
- Batsis, J. A., Mackenzie, T. A., Bartels, S. J., Sahakyan, K. R., Somers, V. K., & Lopez-Jimenez, F. (2016). Diagnostic accuracy of body mass index to identify obesity in older adults: NHANES 1999–2004. *International journal of obesity*, 40(5), 761–767.
- Bhurtyal, A., & Adhikari, D. (2022). Temporal trends, socio-economic inequalities in obesity and responses by federal government, Nepal: a systematic review of observational studies, policies, strategies and plans, 2005–2019. *medRxiv*, 2022–03.

Cohen, A. K., Rai, M., Rehkopf, D. H., & Abrams, B. (2013). Educational attainment and obesity: a systematic review. *Obesity reviews*, 14(12), 989–1005.

Correa, M. M., Thume, E., De Oliveira, E. R. A., & Tomasi, E. (2016). Performance of the waist-to-height ratio in identifying obesity and predicting non-communicable diseases in the elderly population: A systematic literature review. *Archives of gerontology and geriatrics*, 65, 174–182.

Egger, G., Swinburn, B., & Islam, F. A. (2012). Economic growth and obesity: an interesting relationship with world-wide implications. *Economics & Human Biology*, 10(2), 147–153.

Garrido-Miguel, M., Cavero-Redondo, I., Álvarez-Bueno, C., Rodríguez-Artalejo, F., Moreno, L. A., Ruiz, J. R., & Martínez-Vizcaíno, V. (2019). Prevalence and trends of overweight and obesity in European children from 1999 to 2016: a systematic review and meta-analysis. *JAMA pediatrics*, 173(10), e192430.

Goh, L. G., Dhaliwal, S. S., Welborn, T. A., Lee, A. H., & Della, P. R. (2014). Anthropometric measurements of general and central obesity and the prediction of cardiovascular disease risk in women: a cross-sectional study. *BMJ open*, 4(2), e004138.

Goran, M. I. (1998). Measurement issues related to studies of childhood obesity: assessment of body composition, body fat distribution, physical activity, and food intake. *Pediatrics*, 101(3 Pt 2), 505–518.

Kanter, R., & Caballero, B. (2012). Global gender disparities in obesity: a review. *Advances in nutrition*, 3(4), 491–498.

Luhar, S., Timæus, I. M., Jones, R., Cunningham, S., Patel, S. A., Kinra, S., ... & Houben, R. (2020). Forecasting the prevalence of overweight and obesity in India to 2040. *PloS one*, 15(2), e0229438.

McLachlan, S., & NCD Risk Factor Collaboration. (2016). Trends in adult body mass index in 200 countries since 1975: pooled analysis of 1,698 population-based measurement studies with 19.2 million participants. *The Lancet*, 387(10026).

Miranda, J. J., Barrientos-Gutierrez, T., Corvalan, C., Hyder, A. A., Lazo-Porras, M., Oni, T., & Wells, J. C. (2019). Understanding the rise of cardiometabolic diseases in low-and middle-income countries. *Nature medicine*, 25(11), 1667–1679.

Misra, A., Chowbey, P., Makkar, B. M., Vikram, N. K., Wasir, J. S., Chadha, D., & Munjal, Y. P. (2009). Consensus statement for diagnosis of obesity, abdominal obesity and the metabolic syndrome for Asian Indians and recommendations for physical activity, medical and surgical management. *Japi*, 57(2), 163–70.

Mistry, S. K., & Puthussery, S. (2015). Risk factors of overweight and obesity in childhood and adolescence in South Asian countries: a systematic review of the evidence. *Public health*, 129(3), 200–209.

Monteiro, C. A., Moura, E. C., Conde, W. L., & Popkin, B. M. (2004). Socioeconomic status and obesity in adult populations of developing countries: a review. *Bulletin of the world health organization*, 82(12), 940–946.

Neuman, M., Kawachi, I., Gortmaker, S., & Subramanian, S. V. (2013). Urban–rural differences in BMI in low–and middle–income countries: the role of socioeconomic status. *The American journal of clinical nutrition*, 97(2), 428–436.

Non-Communicable Diseases Risk Factor Collaboration. (2019). Rising rural body-mass index is the main driver of the global obesity epidemic in adults. *Nature*, 569(7755), 260–4.

Pavela, G., Lewis, D. W., Locher, J., & Allison, D. B. (2016). Socioeconomic status, risk of obesity, and the importance of Albert J. Stunkard. *Current obesity reports*, 5(1), 132–139.

Perissinotto, E., Pisent, C., Sergi, G., Grigoletto, F., Enzi, G., & ILSA Working Group. (2002). Anthropometric measurements in the elderly: age and gender differences. *British Journal of nutrition*, 87(2), 177–186.

Popkin, B. M., & Ng, S. W. (2022). The nutrition transition to a stage of high obesity and noncommunicable disease prevalence dominated by ultra-processed foods is not inevitable. *Obesity reviews*, 23(1), e13366.

Popkin, B. M., Corvalan, C., & Grummer-Strawn, L. M. (2020). Dynamics of the double burden of malnutrition and the changing nutrition reality. *The Lancet*, 395(10217), 65–74.

Popkin, B.M. (2003). The nutrition transition in the developing world. *Development policy review*, 21(5–6), 581–597.

Pries, A. M., Rehman, A. M., Filteau, S., Sharma, N., Upadhyay, A., & Ferguson, E. L. (2019). Unhealthy snack food and beverage consumption is associated with lower dietary adequacy and length-for-age z-scores among 12–23-month-olds in Kathmandu Valley, Nepal. *The Journal of nutrition*, 149(10), 1843–1851.

Ranjani, H., Mehreen, T. S., Pradeepa, R., Anjana, R. M., Garg, R., Anand, K., & Mohan, V. (2016). Epidemiology of childhood overweight & obesity in India: A systematic review. *Indian Journal of Medical Research*, 143(2), 160–174.

Skinner, A. C., Ravanbakht, S. N., Skelton, J. A., Perrin, E. M., & Armstrong, S. C. (2018). Prevalence of obesity and severe obesity in US children, 1999–2016. *Pediatrics*, 141(3), e20173459.

Templin, T., Cravo Oliveira Hashiguchi, T., Thomson, B., Dieleman, J., & Bendavid, E. (2019). The overweight and obesity transition from the wealthy to the poor in low–and middle–income countries: A survey of household data from 103 countries. *PLoS medicine*, 16(11), e1002968.

Wen, C. P., Cheng, T. Y. D., Tsai, S. P., Chan, H. T., Hsu, H. L., Hsu, C. C., & Eriksen, M. P. (2009). Are Asians at greater mortality risks for being overweight than Caucasians? Redefining obesity for Asians. *Public health nutrition*, 12(4), 497–506.

Wen, C. P., Cheng, T. Y. D., Tsai, S. P., Chan, H. T., Hsu, H. L., Hsu, C. C., & Eriksen, M. P. (2009). Are Asians at greater mortality risks for being overweight than Caucasians? Redefining obesity for Asians. *Public health nutrition*, 12(4), 497–506.

World Cancer Research Fund/American Institute for Cancer Research. (2018). Diet, nutrition, physical activity and cancer: A global perspective (Summary of the third expert report). London: WCRF.

Out-of-pocket health expenditure: A repeated cross-sectional analysis of National Sample Survey data, 2004–11 to 2022

Abhishek Kumar [#]

Several ongoing reforms in India in the health sector are aimed at reducing the financial burdens of the households. Against this backdrop the study presents the estimates of out-of-pocket expenditure and incidence of catastrophic health expenditure.

We analyze out-of-pocket and catastrophic health expenditures using data from three National Sample Survey rounds (2004–05, 2011–12, 2022). Additionally, we report institutional and non-institutional medical expenditure across categories like medicines and diagnostic tests. Using logistic regression, we assess the socioeconomic factors influencing catastrophic health expenditure within a multivariate framework.

The share of total medical expenditure as a percentage of MPCE was 5.9% in 2004–05, and 6.6% in 2022. The incidence of catastrophic health expenditure, if it exceeds 10% of MPCE, was 14.6% in 2004–05, and 15.9% in 2022. Among non-institutional health expenditures, pathological tests accounted for a significant portion. Muslim households; SC and OBC households, casual laborers and households in the upper quintiles are more likely to experience catastrophic health expenditure.

This study underscores the urgent need for well-designed policy interventions to reduce OOP expenses. Expanding insurance coverage and providing subsidized treatment at public facilities are key steps toward achieving universal health coverage.

[#] Corresponding Author: Abhishek Kumar, FLAME University, Pune, Maharashtra, India
Email: abhishekkmr48@gmail.com

Out of pocket expenditure refers to the direct payment made by individuals while seeking healthcare at private or public facilities (Wagstaff et al., 2020). The payments are not reimbursable from insurance and government scheme. Out of pocket expenditure can have significant consequences for households. In absence of insurance coverage and rising medical costs, the households are forced to forgo treatment, discontinue treatment or have to settle for substandard quality of services. On the other hand, catastrophic health expenditure occurs when a household out of pocket expenditure exceeds a certain threshold of their income or consumption and can lead to financial distress (Wagstaff et al., 2020). Catastrophic health expenditure is an important indicator for assessing the financial protection in health care systems.

To ensure the well-being of people, the Sustainable Development Goals which is a collective commitment by many countries, emphasize the need for providing universal health coverage (Royston et al., 2020). UHC includes providing equitable access to health care services to everyone. India as a signatory to the 2030 agenda for sustainable development has taken measure to achieve the targets laid out in SDG framework and ensure UHC for its population. However, out of pocket health expenditure is a major cause of concern for India as the coverage of insurance is low and the demographics are changing substantially over time (Khan et al., 2021; Dang et al., 2021; Sarwal & Kumar, 2021).

Here it is worth noting that Out-of-pocket (OOP) expenditure can be measured in several ways to assess its financial burden on households (Wagstaff et al., 2020). Absolute measures include total OOP expenditure, which captures the direct amount spent on healthcare services, and per capita OOP expenditure, which calculates the average spending per person (Wagstaff et al., 2020; Banthin et al., 2008). The relative measures express OOP expenditure as a proportion of total household expenditure or income, providing insight into financial strain (Wagstaff et al., 2020). Another critical measure is catastrophic health expenditure (CHE), which occurs when OOP spending exceeds a predefined threshold, such as 10% of total household expenditure, or 40% of a household's non consumption expenditure (Wagstaff et al., 2020; Berki, 1986). Additionally, OOP spending can lead to impoverishment, where households fall below the poverty line after paying for healthcare (Wagstaff & Doorslaer, 1992; Wagstaff & Doorslaer, 2003). The poverty headcount approach quantifies this by measuring the proportion of households pushed into poverty due to medical expenses.

Measurement of out-of-pocket is challenging given the different nature of diseases and recall biases. Estimates are usually based on household survey data and consumption expenditure is used as the denominator in absence of reliable estimates for income. Relative measures are usually used to allow temporal comparisons. There are several studies which have tried to estimate the out-of-pocket expenditure for India based on household survey data. Some studies found

that CHE has increased between 1993 and 2011 based on consumption expenditure round of NSS (Karan et al., 2014; Selvaraj et al., 2018). A recent study based on NSS data also found that OOP expenditure was catastrophic for 17% of the households (Selvaraj et al., 2018). In fact, a comparison for countries in South Asia reveals that India ranks third among countries which report high OOP expenditure. As per national health accounts data, the out-of-pocket payments for India are estimated to be 39% of the total health expenditure (NHSRC, 2024).

In addition, the overall OOP payments conceal the expenditure for different diseases. For instance, the OOP expenditure for cancer and other NCDs is quite high and more detrimental and impoverishing (Dhankar et al., 2021; Yadav et al., 2021). However, the problem with these studies is the small sample size and inadequate representation in the overall sample. There are other studies which have estimated OOP expenditure for certain age groups and established association with socio-economic characteristics. For instance, catastrophic expenditure is commonest among households comprising only people aged 60 years or older (Yadav et al., 2021). There are few studies which have also estimated the expenditure on components of health such as medicine and pathological tests (Selvaraj et al., 2018; Prinja et al., 2022).

There is a consensus that increasing public spending could lead to a reduction in OOP expenditure (WHO, 2021). Although, India has in place the Ayushman Bharat yojana, but the spending as a share of GDP on public health spending has remained low over the past few decades (Jakovljevic et al., 2022). Over, the past few years, the medical inflation has been very high, the doctor consultation and diagnostic charge has also increased many folds. The current insurance coverage across India shows a divide (Keshri & Ghosh, 2019). The states in south have better coverage. Although, the National Health policy strives to mitigate the impact of the high OOP payments but it is not visible in the budget. An analysis of the OOP expenditure on various components could be a good starting point to provide insights for effective policy making. These estimates are essential for informed decision-making on fund allocation and drafting health packages. Therefore, we present detailed estimates of health-related costs and their percentage contributions using the latest (2022) round of the National Sample Survey on Consumption Expenditure, which captures the post-COVID-19 period and allows for assessment of how the pandemic influenced household health spending patterns.

Data and Methods

- Data

The study is based on data from three rounds of Consumer expenditure survey conducted during 2004-05, 2011-12 and 2022 by the National Sample Survey

Organization (NSSO). A key focus of these surveys is to collect data on the socio-economic and demographic characteristics of the households. Information is collected on food as well as non-food items. The primary objective of the survey is to estimate and analyze the distribution of households' Monthly Per Capita Expenditure (MPCE) across rural and urban areas.

The information on out-of-pocket expenditure incurred by household on hospitalization (institutional expenditure) during last 365 days and non-hospitalization (non-institutional expenditure) for last 30 days was collected separately during the survey. Information on different items for Medicine, Doctor's/surgeon's fee, X-ray, ECG, pathological test, etc.; hospital & nursing home charges; and other medical expenses was collected for both hospitalization and non-hospitalization cases.

- Survey Design

The Consumer expenditure survey follows a multi-stage survey design covering the whole of India. Clusters are formed by grouping districts with similar characteristics, from which urban and rural areas are identified for data collection. In the first stage, the selection process follows the probability proportional to size (PPS) method. In other words, this design involves dividing the population into first stage units (FSUs) and ultimate stage units (USUs). The first stage units (FSUs) are villages, blocks, or sub-units (SUs). The ultimate stage units (USUs) are households which are circular systematically selected to constitute the final sample.

A total of 1,24,644 households were interviewed in 2004-05; 1,01,662 (in type 1) and 1,01,651 (in type 2) during 2011-12 and 2,61,746 during 2022. In the 2011-12 round two different data collection schedules were used- "Type 1" and "Type 2". The difference between the two is that Type 1 generally uses a longer recall period (last 30 days and last 365 days) compared to Type 2 which uses a shorter recall period (last 7 days) for specific food items. We have used data for both the schedules for the analysis.

• Outcomes

The data on health expenditure was used to create three new variables a) per capita medical expenditure (institutional and non-institutional) b) OOP spending on medical as well as medicine as a share of MPCE and non-food MPCE c) percentage of households reporting catastrophic health expenditure. Two thresholds were established for catastrophic health expenditure: one where medical expense exceeded 10% of Monthly Per Capita Expenditure (MPCE) and another where medical expenses exceeded 40% of non-food MPCE. The estimates are also provided separately for rural and urban areas. The break-up of the medical expenditure for different rounds is also provided.

• Independent variables

Socio-economic differentials can have a significant impact on the incidence of OOP expenditure. In this study, we considered religion, social group, household type, household size and quintiles based on MPCE as the key determinants. Social group was categorized as scheduled tribes (ST), scheduled castes (SC), other backward classes (OBC) and other castes. In addition, the religion (Hindu, Muslim and Others) and household type based on employment type (Self-employed, Salaried, Casual Labour and Not employed) were considered. The household were categorized in five quintiles based on value of MPCE.

• Statistical Analysis

We report medical expenditure for both institutional and non-institutional sources across various survey rounds and examine expenditure inequalities among different social groups. All household expenditure data for 2004, 2011, and 2023 were converted into real terms (constant 2011 prices) using the Consumer Price Index (CPI). Additionally, we estimate the number of households experiencing catastrophic health expenditure. To analyse the factors associated with catastrophic health expenditure, we use logistic regression within a multivariate framework, adjusting for socioeconomic status (SES) factors. The results are reported as Odds Ratios (OR) with corresponding 95% confidence intervals. These odds ratios provide a relative measure of effect, enabling comparisons between groups relative to a reference group. The analysis was conducted using Stata 12.

Results

• Socio-economic Characteristics of the households

The distribution of household socio-economic characteristics has remained largely consistent across survey rounds (Table 1). In 2022, 81.5% of the interviewed households were Hindu, 44.7% belonged to the OBC category, and 57.1% were self-employed. The percentage of self-employed households increased from 52.6% in 2004 to 57.1% in 2022, while the share of casual labour households declined from 29.3% in 2004 to 23% in 2022.

• Financial Burden on the households

First, we present estimates of Monthly Per Capita Expenditure (MPCE), total medical expenditure, and expenditure on medicines (both institutional and non-institutional) for each survey round (Table 2). Household MPCE increased from ₹1223.7 in 2004 to ₹2405.5 in 2022, while non-food MPCE rose from ₹634.3 in 2004-05 to ₹1359.9 in 2022. The mean per capita institutional and non-institutional medical expenditures were ₹52.4 and ₹107.2 in 2022, respectively. In

Table 1: Sample Characteristics, NSS 2004, 2011–12 (Type 1 and 2) and 2022

	2004		2011 (Type 1)		2011 (Type 2)		2022	
	N	%	No.	%	No.	%	No.	%
Religion								
Hindu	102402	82.2	82847	81.5	82769	81.4	213249	81.5
Muslim	15774	12.7	13869	13.6	13833	13.6	34931	13.3
Others	6468	5.2	4947	4.9	5049	5	13566	5.2
Total	124644	100	101662	100	101651	100	261746	100
Social Group								
ST	10761	8.6	9079	8.9	9150	9	26291	10
SC	24413	19.6	19351	19	19412	19.1	51802	19.8
OBC	51031	40.9	44790	44.1	44240	43.5	116908	44.7
Others	38440	30.8	28442	28	28849	28.4	66746	25.5
Total	124644	100	101662	100	101651	100	261746	100
Household type								
Self-employed	65616	52.6	51831	51	51248	50.4	149417	57.1
Salaried	12438	10	17724	17.4	17468	17.2	37761	14.4
Casual Labour	36536	29.3	27850	27.4	28426	28	60295	23
Not employed	10055	8.1	4230	4.2	4476	4.4	14273	5.5
Total	124644	100	101635	100	101618	100	261746	100
MPCE based Quintiles								
Quintile 1	24934	20	20339	20	20331	20	52350	20
Quintile 2	24931	20	20329	20	20331	20	52350	20
Quintile 3	24922	20	20331	20	20332	20	52353	20
Quintile 4	24931	20	20333	20	20330	20	52345	20
Quintile 5	24926	20	20330	20	20327	20	52348	20
Total	124644	100	101662	100	101651	100	261746	100

2022, institutional expenditure on medicines was ₹21.1, while non-institutional expenditure on medicines was at ₹82.5. The share of total medical expenditure (both institutional and non-institutional) as a percentage of MPCE was 5.9% in 2004–05, 6.8% in 2011 (Type 1), and 6.6% in 2022, showing a decline between 2011 and 2022. The share of medicine expenditure decreased from 4.53% in 2011 to 4.31% in 2022. The total medical expenditure in rural areas (7.1%) is higher as compared to urban areas (5.9%) (Table S1).

• Catastrophic Health Expenditure

The incidence of catastrophic health expenditure, based on the first approach, where health expenditure is considered catastrophic if it exceeds 10% of MPCE,

Table 2: Financial burden indicators, India, 2004–2005, 2011–2012 and 2022–23

NSS round	2004	2011 (Type-1)	2011 (Type -2)	2022
Monthly Per Capita Expenditure (Current price)				
Household MPCE	1223.7	1627.1	1772.7	2405.5
	[1217.7,1229.7]	[1617.4,1636.9]	[1761.6,1783.7]	[2398.5,2412.4]
Household Non-food MPCE	634.3	919.2	912.1	1359.9
	[629.7,639]	[911.3, 927.1]	[903.8, 920.3]	[1354.6,1365.1]
Monthly Total medical expenditure per capita				
Medical expenditure institutional				
365 days (converted to 30 days)	19.0	36.4	36.7	52.4
	[18.1,19.9]	[34.8,37.9]	[35.2,38.2]	[51.2,53.6]
Medical non institutional				
30 days	52.6	74.3	72.9	107.2
	[51.8,53.4]	[73.2,75.5]	[71.8,74.1]	[106.2,108.1]
Monthly Expenditure on Medicines per capita				
Expenditure on medicine (institutional)	7.8	15.5	14.6	21.1
	[1.8,2.1]	[14.5,16.5]	[14,15.3]	[20.6,21.7]
Expenditure on medicine (non-institutional)	43.3	58.2	56.8	82.5
	[42.6,43.9]	[57.3,59]	[56.1,57.6]	[81.8,83.1]
Share in total MPCE				
Total medical as Share of MPCE (%)	5.9	6.8	6.2	6.6
	[5.8,5.9]	[6.7,6.9]	[6.1,6.3]	[6.6,6.7]
Total medicine as Share of MPCE (%)	4.2	4.5	4	4.3
	[4.1,4.2]	[4.5,4.6]	[4,4.1]	[4.3,4.3]
Share in total non-food MPCE				
Total medical as Share of non-food MPCE (%)	11.3	12	12	11.7
	[11.2,11.4]	[11.9,12.2]	[11.9,12.1]	[11.7,11.8]
Total medicine as Share of non-food MPCE (%)	8.0	8	7.8	7.6
	[8,8.1]	[7.9,8.1]	[7.8,7.9]	[7.6,7.7]

was 14.6% in 2004–05, 17.3% in 2011 (Type 1), and 15.9% in 2022 (Table 4). The percentage of households reporting catastrophic expenditure on medicines under this approach was 10.1% in 2004–05, 10.4% in 2011 (Type 1), and 9.1% in 2022. Using the second approach, where health expenditure is considered catastrophic if it exceeds 40% of non-food MPCE, the incidence was 4.6% in 2011 (Type 1) and 4.1% in 2022. The incidence was 4.1% in rural areas in 2022 (Table S2).

Table 3: Percentage of households incurring catastrophic payments with respect to total OOP spending and medicines OOP spending, India, 2004–2005, 2011–2012 and 2022–23

	2004	2011 (Type-1)	2011 (Type -2)	2022
Catastrophe using MPCE				
Percentage of Households reporting total medical expenditure >10%	14.6	17.3	15.3	15.9
	[14.4,14.8]	[17.1,17.6]	[15.1,15.5]	[15.8,16]
Percentage of Households reporting total expenditure on medicines >10%	10.1	10.4	9.4	9.1
	[10,10.3]	[10.2,10.6]	[9.2,9.5]	[9,9.2]
Catastrophe using Non-food MPCE				
Percentage of Households reporting total medical expenditure >40%	4.5	4.6	5.0	4.1
	[4.4,4.6]	[4.5,4.8]	[4.9,5.1]	[4,4.1]
Percentage of Households reporting total expenditure on medicines >40%	2.0	1.7	1.8	1.2
	[1.9,2.1]	[1.6,1.8]	[1.7,1.9]	[1.1,1.2]

• Expenditure on Medical items

The monthly per capita expenditure (MPCE) on various items shows that spending on medicines remained the highest, reaching ₹21.1 in 2022, up from ₹7.8 in 2004 and ₹15.5 in 2011 (Type 1). The share of institutional medicine expenditure in MPCE increased from 0.6% in 2004 to 1% in 2011, and was at 0.9% in 2022. In contrast, the share of non-institutional medicine expenditure decreased from 3.5% in 2004 to 3.4% in 2022. Among non-institutional health expenditures, pathological tests accounted for a significant portion. In 2022, the monthly per capita expenditure on non-institutional health items was ₹82.5.

• Catastrophic expenditure by household characteristics

The percentage of households experiencing catastrophic health expenditure, based on Approach 1 (where health expenditure is considered catastrophic if it exceeds 10% of MPCE), was 17.4% in 2011 (type 1) and 15.7% in 2022 among Hindu households. Among Scheduled Caste (SC) households, the incidence declined from

18.5% in 2011 (type 1) to 15.4% in 2022, while for OBC households, it dropped from 18.4% to 16.3% over the same period. There were no significant changes in the incidence of catastrophic health expenditure among salaried and non-employed households. By economic status, the incidence declined from 10.8% to 8.6% among households in the bottom quintile and from 17.3% to 15.0% among those in the top quintile between 2011 (type 1) and 2022.

Table 5: Percentage households who incurred catastrophic expenditure (as share of total MPCE) by background characteristics

Background Characteristics	2004		2011 Type 1		2011 Type-2		2022	
	N	%	No.	%	No.	%	No.	%
Religion								
Hindu	102402	14.5	82847	17.4	82769	15.3	213249	15.7
Muslim	15774	15	13869	16.7	13833	15.3	34931	16.4
Others	6468	14.8	4947	18	5049	15.8	13566	17.8
Total	124644	14.6	101662	17.3	101651	15.3	261746	15.9
Social Group								
ST	10761	8.9	9079	9.9	9150	9.8	26291	10
SC	24413	14.4	19351	18.5	19412	15.9	51802	15.4
OBC	51031	15.7	44790	18.4	44240	16.4	116908	16.3
Others	38440	14.7	28442	17.2	28849	15	66746	17.9
Total	124644	14.6	101662	17.3	101651	15.3	261746	15.9
Household type								
Self-employed	65616	14.4	51831	16.9	51248	15.3	149417	15.4
Salaried	12438	12.3	17724	15.6	17468	14.1	37761	15
Casual Labour	36536	15.1	27850	18	28426	15	60295	15.8
Not employed	10055	16.2	4230	25	4476	22.5	14273	24.4
Total	124644	14.6	101635	17.3	101618	15.3	261746	15.9
MPCE Quintiles								
Quintile 1	24934	7.4	20339	10.8	20331	8.9	52350	8.6
Quintile 2	24931	10.9	20329	13.4	20331	12.4	52350	12.6
Quintile 3	24922	13.2	20331	16.4	20332	14.8	52353	15.3
Quintile 4	24931	17.3	20333	19.8	20330	17.1	52345	18.5
Quintile 5	24926	23.9	20330	26.2	20327	23.4	52348	24.5
Total	124644	14.6	101662	17.3	101651	15.3	261746	15.9

• Association of Catastrophic expenditure with socio-economic characteristics of the household

Casual laborers are 20% [OR-1.20, 95% CI: 1.16-1.24] more likely to incur catastrophic health expenditure, while non-employed individuals also tend to have higher health expenditures. Households in the upper quintiles are three times more likely [OR-3.71, 95% CI: 3.51-3.92] to experience catastrophic health expenditure.

SC [OR=1.47, 95% CI: 1.38-1.57] and OBC [OR=1.47, 95% CI: 1.39-1.56] households face a 47% higher likelihood [OR=1.47, 95% CI: 1.38-1.57] of experiencing such expenses. Muslim households are 8% [OR=1.08, 95% CI: 1.03-1.13] more likely to incur catastrophic health expenditure compared to Hindu households.

Table 6: Logistic regression results for experience of catastrophic health expenditure

		2004	2011 Type 1	2011 Type-2	2022
Religion	Hindu	1	1	1	1
		[1.00,1.00]	[1.00,1.00]	[1.00,1.00]	[1.00,1.00]
	Muslim	1.20***	1.03	1.07	1.08***
		[1.10,1.31]	[0.93,1.15]	[0.96,1.20]	[1.03,1.13]
	Others	0.83***	0.91	0.9	1.02
		[0.75,0.91]	[0.81,1.03]	[0.79,1.03]	[0.96,1.09]
Social group	ST	1	1	1	1
		[1.00,1.00]	[1.00,1.00]	[1.00,1.00]	[1.00,1.00]
	SC	1.58***	1.88***	1.62***	1.47***
		[1.40,1.79]	[1.61,2.20]	[1.36,1.93]	[1.38,1.57]
	OBC	1.59***	1.76***	1.54***	1.47***
		[1.41,1.79]	[1.51,2.04]	[1.31,1.81]	[1.39,1.56]
	Others	1.24***	1.46***	1.24**	1.52***
		[1.10,1.40]	[1.25,1.70]	[1.05,1.46]	[1.44,1.62]
Household Type	Self-employed	1	1	1	1
		[1.00,1.00]	[1.00,1.00]	[1.00,1.00]	[1.00,1.00]
	Salaried	0.83***	0.80***	0.81***	0.97
		[0.75,0.91]	[0.73,0.87]	[0.73,0.88]	[0.93,1.02]
	Casual Labour	1.45***	1.30***	1.14***	1.20***
		[1.36,1.56]	[1.19,1.42]	[1.04,1.25]	[1.16,1.24]
	Not employed	0.95	1.54***	1.51***	1.72***
		[0.87,1.04]	[1.37,1.74]	[1.33,1.71]	[1.64,1.80]

		2004	2011 Type 1	2011 Type-2	2022
MPCE Quintile	Quintile 1	1 [1.00,1.00]	1 [1.00,1.00]	1 [1.00,1.00]	1 [1.00,1.00]
	Quintile 2	1.62*** [1.45,1.82]	1.32*** [1.14,1.53]	1.50*** [1.29,1.75]	1.54*** [1.45,1.63]
	Quintile 3	2.15*** [1.92,2.40]	1.76*** [1.53,2.02]	1.92*** [1.65,2.22]	1.96*** [1.85,2.07]
	Quintile 4	3.19*** [2.86,3.55]	2.34*** [2.04,2.67]	2.36*** [2.05,2.72]	2.53*** [2.39,2.67]
	Quintile 5	5.36*** [4.82,5.96]	3.62*** [3.17,4.14]	3.74*** [3.25,4.31]	3.71*** [3.51,3.92]
	Constant	0.04*** [0.03,0.04]	0.06*** [0.05,0.07]	0.06*** [0.05,0.07]	0.05*** [0.05,0.05]
	N	124644	101635	101618	261746

Discussion

We estimate the out-of-pocket expenditure and incidence of catastrophic health expenditure using two commonly used approaches based on data from three rounds of National sample survey on consumption expenditure. Following are the four salient findings from this study which could provide insights into financial burden on households.

First, the share of monthly per capita expenditure (MPCE) on both hospitalization and non-hospitalization in total and non-food MPCE has declined over time. A study based on consumer expenditure surveys from 1993–94 to 2011–12 reported an increase in the share of medical expenses in MPCE from 4.84% to 6.77% (Selvaraj et al., 2018). While our study indicates a decline between 2011 and 2022, the magnitude of reversal is not as large. However, these findings are in line with other studies based on health consumption and morbidity survey data which show a decrease in the out-of-pocket (OOP) expenditure share between the 71st (2014) and 75th (2017–18) rounds of the NSS (Mohanty & Dwivedi, 2021). Furthermore, National Health Accounts data reveal a decline in OOP expenditure as a share of total health expenditure, from 64.2% in 2013–14 to 39.4% in 2021–22, a positive trend (NHSRC, 2024). This shift may be attributed to increased government investment in public healthcare initiatives like Ayushman Bharat, improving access to affordable healthcare and reducing direct medical expenses (Parmar et al., 2023). Additionally,

a rise in government health expenditure as a percentage of GDP could be another factor (NHSRC, 2024). Some studies suggest that countries with an increase in national health expenditure as a share of GDP incur lower expenditure on OOP (WHO, 2021). However, it should be noted that several health insurance schemes such as the Rajiv Aarogyasri Health Insurance Scheme in Andhra Pradesh, the Rajiv Gandhi Jeevandayee Arogya Yojana in Maharashtra, and the centrally implemented Rashtriya Swasthya Bima Yojana (RSBY), were introduced to reduce financial hardships for the poor. However, they did not perform as expected (Sriram & Khan, 2020; Prinja et al., 2017).

Second, the percentage of household reporting CHE based on approach one (greater than 10% as a share of MPCE) has declined from 17.3% to 15.9% between 2011 and 2022 while based on approach 2, greater than 40% of non-food MPCE) has declined from 4.6 to 4.1%. Interestingly, an earlier study had showed that the percentage of households reporting CHE had increased between 1993-94 and 2011-12 from 13.9 to 17.9% based on approach 1 but had actually declined from 9.7 to 4.9% based on approach two (Selvaraj et al., 2018). Study based on health consumption and morbidity survey data also show that the incidence of CHE has declined from 23.45 in 2013-14 to 16.69 in 2017-18 (Mohanty & Dwivedi, 2021). While recent studies based on 75th round has reported that a total of 20.4% households faced CHE for hospitalization based on the average per capita consumption expenditure while 28% of households were grappling with the complexity of financial burden due to elevated inpatient healthcare (Yadav et al., 2021; Panda et al., 2024).

Third, we found that the share of household expenditure on medicines has remained largely unchanged. Most of this spending goes toward medicines and hospital or nursing charges for hospitalization, as well as medicines and pathological tests for non-hospitalization cases (Selvaraj et al., 2018; Ambade et al., 2022). A previous study estimated that spending on medicines accounts for approximately 7.6% of non-food MPCE, indicating that the financial burden of medical expenses has not significantly changed (Selvaraj et al., 2018). The National Health Policy 2017 emphasized the need to provide free medicines in public health facilities by increasing funding and improving drug procurement and supply chain mechanisms (MoHFW, 2017). States like Tamil Nadu and Rajasthan have well-functioning supply chain systems that ensure drug availability (Selvaraj et al., 2014; Kokilam et al., 2016). Additionally, the Indian government's recent initiative to exempt basic duty on cancer drugs, as well as for rare and chronic diseases, is a positive step. However, it remains to be seen whether these benefits will effectively reach patients.

Lastly, we found that out-of-pocket (OOP) expenditure varies based on households' socio-economic characteristics. Specifically, households that are Muslim, belong to the Other Backward Classes (OBC), work as casual laborers, or are in the upper wealth quintiles are more likely to incur catastrophic health expenditure (CHE). Several studies have highlighted the higher incidence of CHE among certain groups

which raises concerns about the effectiveness of existing health schemes (Karan et al., 2014; Pandey et al., 2018; Gaddam & Rao, 2023). One possible reason for the higher incidence of catastrophic health expenditure (CHE) among the richest quintile is their greater capacity and willingness to seek and pay for healthcare, including costlier institutional care and specialized treatments. In contrast, poorer households often delay or forgo medical treatment altogether because of financial barriers, limited access to quality healthcare, or reliance on public facilities with lower out-of-pocket costs. As a result, while the poor may face unmet health needs, the rich appear to incur higher medical expenses, leading to a relatively greater incidence of CHE despite their stronger financial position.

Limitations

This study has several limitations. First, the household survey data relies on self-reported expenditures, which may be affected by recall bias. Since there is no way to validate health expenditures from other sources, the data's accuracy remains uncertain. Additionally, households may underreport expenses due to social desirability bias. Second, the analysis only accounts for direct costs, excluding indirect costs such as lost income, transportation, and accommodation expenses. As a result, catastrophic health expenditures may be underestimated. Third, the study focuses on households that had the ability to pay for treatment, excluding those who did not seek medical care due to financial constraints. This might have underestimated the proportion of households experiencing catastrophic health expenditures.

Furthermore, the study does not include a disaggregated analysis by disease type. When CHE is assessed only at the household level, high costs associated with specific illnesses such as chronic conditions, maternal care, or hospitalization may be diluted by averaging total household spending. As a result, disease-specific or service-specific analyses can reveal more precise patterns of financial hardship, particularly for conditions that disproportionately affect certain demographic or socioeconomic groups. Therefore, disaggregated analyses are required to better inform targeted health financing and protection policies (Ataguba et al., 2024).

Finally, the use of cross-sectional survey data introduces temporal limitations. A longitudinal database is needed to capture the true burden of catastrophic health expenditures over time, especially to account for the impact of policy changes.

Conclusion

This study offers several policy-relevant insights. The findings suggest that out-of-pocket medical spending continues to be a major source of financial burden for households, with outpatient care, particularly diagnostic tests playing a significant

role. Vulnerability to catastrophic health expenditure remains pronounced among socially and economically disadvantaged groups as well as higher-income households who tend to seek more expensive care. These patterns reinforce the need for stronger financial protection mechanisms, including targeted subsidies for outpatient services and improved access to affordable diagnostics and medicines. Strengthening public health infrastructure and ensuring the consistent availability of essential drugs, as emphasized in the National Health Policy 2017, remain critical to reducing household financial strain and advancing toward universal health coverage.

References

- Ambade, M., Sarwal, R., Mor, N., Kim, R., & Subramanian, S. V. (2022). Components of out-of-pocket expenditure and their relative contribution to economic burden of diseases in India. *JAMA Network Open*, 5(5), e2210040–e2210040.
- Ataguba, J. E., Ichoku, H. E., Ingabire, M. G., & Akazili, J. (2024). Financial protection in health revisited: Is catastrophic health spending underestimated for service-or disease-specific analysis?. *Health Economics*, 33(6), 1229–1240.
- Banthin, J. S., Cunningham, P., & Bernard, D. M. (2008). Financial burden of health care, 2001–2004. *Health Affairs*, 27(1), 188–195.
- Berki, S. E. (1986). A look at catastrophic medical expenses and the poor. *Health affairs*, 5(4), 138–145.
- Dang, A., Dang, D., & Vallish, B. N. (2021). Importance of evidence-based health insurance reimbursement and health technology assessment for achieving universal health coverage and improved access to health in India. *Value in Health Regional Issues*, 24, 24–30.
- Dhankhar, A., Kumari, R., & Bahurupi, Y. A. (2021). Out-of-pocket, catastrophic health expenditure and distress financing on non-communicable diseases in India: A systematic review with meta-analysis. *Asian Pacific Journal of Cancer Prevention: APJCP*, 22(3), 671.
- Gaddam, R., & Rao, K. R. (2023). Incidence, inequality and determinants of catastrophic health expenditure in India. *Journal of Health Management*, 25(1), 30–39.
- Jakovljevic, M., Lamnissos, D., Westerman, R., Chattu, V. K., & Cerda, A. (2022). Future health spending forecast in leading emerging BRICS markets in 2030: health policy implications. *Health research policy and systems*, 20(1), 23.
- Karan, A., Selvaraj, S., & Mahal, A. (2014). Moving to universal coverage? Trends in the burden of out-of-pocket payments for health care across social groups in India, 1999–2000 to 2011–12. *PloS one*, 9(8), e105162.
- Keshri, V. R., & Ghosh, S. (2019). Health Insurance for Universal Health Coverage in India: A critical analysis based on coverage, distribution and predictors from National Family Health Survey–4 Data.

Khan, P. K., Perkins, J. M., Kim, R., Mohanty, S. K., & Subramanian, S. V. (2021). Multilevel population and socioeconomic variation in health insurance coverage in India. *Tropical Medicine & International Health*, 26(10), 1285–1295.

Kokilam, M. B., Joshi, H. G., & Kamath, V. G. (2016). Strengthening the pharmaceutical supply chain management with information communication technology intervention: a windfall to the Indian Rural Public Healthcare System. *Journal of Health Management*, 18(2), 274–289.

Ministry of Health and Family Welfare (MoHFW) GoI. National Health Policy, 2017. 2017. https://www.nhp.gov.in/NHPfiles/national_health_policy_2017.pdf

Mohanty, S. K., & Dwivedi, L. K. (2021). Addressing data and methodological limitations in estimating catastrophic health spending and impoverishment in India, 2004–18. *International Journal for Equity in Health*, 20, 1–18.

National Health Systems Resource Centre (2024). National Health Accounts Estimates for India (2021–22). New Delhi: Ministry of Health and Family Welfare, Government of India. <https://nhsrcindia.org/sites/default/files/2024-09/NHA%202021-22.pdf>

NSSO. “Level and Pattern of Consumer Expenditure, 2004–05”, Ministry of Statistics and Programme Implementation, Government of India: 2006.

NSSO. “Level and Pattern of Consumer Expenditure, 2022–23”, Ministry of Statistics and Programme Implementation, Government of India: 2024.

NSSO. Level and Pattern of Consumer Expenditure, 2011–12, Ministry of Statistics and Program Implementation. 2014.

Panda, H. S., Rout, H. S., & Jakovljevic, M. (2024). Catastrophic health expenditure of inpatients in emerging economies: evidence from the Indian subcontinent. *Health Research Policy and Systems*, 22(1), 104.

Pandey, A., Kumar, G. A., Dandona, R., & Dandona, L. (2018). Variations in catastrophic health expenditure across the states of India: 2004 to 2014. *PLoS One*, 13(10), e0205510.

Parmar, D., Strupat, C., Srivastava, S., Brenner, S., Parisi, D., Ziegler, S., ... & De Allegri, M. (2023). Effects of the Indian National Health Insurance Scheme (PM-JAY) on hospitalizations, out-of-pocket expenditures and Catastrophic expenditures. *Health Systems & Reform*, 9(1), 2227430.

Prinja, S., Chauhan, A. S., Karan, A., Kaur, G., & Kumar, R. (2017). Impact of publicly financed health insurance schemes on healthcare utilization and financial risk protection in India: a systematic review. *PloS one*, 12(2), e0170996.

Prinja, S., Kumar, S., Sharma, A., Kar, S. S., Tripathi, N., Dumka, N., ... & Aggarwal, A. K. (2022). What is the out-of-pocket expenditure on medicines in India? An empirical assessment using a novel methodology. *Health policy and planning*, 37(9), 1116–1128.

Royston, G., Pakenham-Walsh, N., & Zielinski, C. (2020). Universal access to essential health information: accelerating progress towards universal health coverage and other SDG health targets. *BMJ Global Health*, 5(5), e002475.

Sarwal, R., & Kumar, A. (2021). Health insurance for India's missing middle.

Selvaraj, S., Farooqui, H. H., & Karan, A. (2018). Quantifying the financial burden of households' out-of-pocket payments on medicines in India: a repeated cross-sectional analysis of National Sample Survey data, 1994–2014. *BMJ open*, 8(5), e018020.

Selvaraj, S., Mukhopadhyay, I., Kumar, P., Aisola, M., Datta, P., Bhat, P., ... & Pachauli, C. (2014). Universal access to medicines: evidence from Rajasthan, India. *WHO South-East Asia journal of public health*, 3(3–4), 289–299.

Sriram, S., & Khan, M. M. (2020). Effect of health insurance program for the poor on out-of-pocket inpatient care cost in India: evidence from a nationally representative cross-sectional survey. *BMC Health Services Research*, 20, 1–21.

Wagstaff, A., & Doorslaer, E. V. (2003). Catastrophe and impoverishment in paying for health care: with applications to Vietnam 1993–1998. *Health economics*, 12(11), 921–933.

Wagstaff, A., & Van Doorslaer, E. (1992). Equity in the finance of health care: some international comparisons. *Journal of health economics*, 11(4), 361–387.

Wagstaff, A., Eozenou, P., & Smitz, M. (2020). Out-of-pocket expenditures on health: a global stocktake. *The World Bank Research Observer*, 35(2), 123–157.

World Health Organization. (2021). Global expenditure on health: public spending on the rise?. In *Global expenditure on health: public spending on the rise?*.

Yadav, J., Menon, G. R., & John, D. (2021). Disease-specific out-of-pocket payments, catastrophic health expenditure and impoverishment effects in India: an analysis of National Health Survey data. *Applied Health Economics and Health Policy*, 19(5), 769–782.

Yadav, J., Allarakha, S., Menon, G. R., John, D., & Nair, S. (2021). Socioeconomic impact of hospitalization expenditure for treatment of noncommunicable diseases in India: a repeated cross-sectional analysis of national sample survey data, 2004 to 2018. *Value in Health Regional Issues*, 24, 199–213..

Livelihood and Status of Tobacco Processing Workers: Insights from Selected States in India

Nayanatara S.Nayak*, Rudra N. Mishra#, Karabi Mujumdar+,
Tara Nair^, N.L.Narasimha Reddy\$

Processing is one of the prime activities in tobacco production and is a labour-intensive activity in India. It involves processes starting from cleaning raw tobacco leaves to crushing them into different sizes/particles or making fine dust in accordance with the requirements of the industries producing bidi, cigarette, zarda, mouth freshener, pan masala, khaini, snuff, and chewing tobacco. There is also a lack of information on the number of processing workers, their livelihood, and, the health hazards of working in tobacco processing units in India. This paper discusses the working conditions and livelihoods of tobacco processing workers and their health status. The study followed snowball sampling to identify a sample of 500 processing workers spread out in the states of Andhra Pradesh, Gujarat, Karnataka, Maharashtra, and West Bengal. The poor working conditions, lack of social security benefits, low standard of living among a majority of workers, reporting of chronic illness among processing workers, absence of scientific studies on the health conditions of workers, and a majority of processing workers not being happy with their employment are the reasons enough for policy intervention to look into their working conditions and find out ways and means to rehabilitate them in other occupations and wean away the new entrants into tobacco processing job market. There is a need for scientific studies based on a large sample of processing workers capturing both clinically diagnosed and self-reported health symptoms for validating the linkage between exposure to tobacco dust and health problems among workers.

Keywords: Tobacco, Processing, Livelihood, Workers

*Corresponding Author: Rudra N. Mishra, Gujarat Institute of Development Research, Ahmedabad, Gujarat, India
Email: rudraam@gmail.com

Centre for Multi-disciplinary Development Research, Dharwad, Karnataka, India

^ Gujarat Institute of Development Research, Ahmedabad, Gujarat, India

+ Freelance Consultant, New Delhi, India

\$ Poverty Learning Foundation, Hyderabad, Andhra Pradesh, India

The final product of tobacco before reaching the consumer involves activities ranging from farming, harvesting, curing, processing, grading, packaging, marketing, manufacturing, distribution, and retailing. Processing is one of the prime activities in tobacco production and is a labour-intensive activity in India. It involves processes starting from cleaning raw tobacco leaves to crushing them into different sizes or fine dust. The crushed leaves differ in size and substance in accordance with the requirements of the industries producing bidi, cigarette, zarda, mouth freshener, pan masala, khaini, snuff, and chewing tobacco. In total it involves processes transforming leaves into powder.

Women are largely involved in processing activities. Tobacco processing is carried out mainly in Karnataka, Gujarat, Andhra Pradesh, West Bengal, and Maharashtra. These are the states, which cultivate different varieties of tobacco.

There is a dearth of information on tobacco processing in India as it is largely in the unorganized sector. Taking into account the production estimates for the entire bidi and chewing tobacco in the country (around 60% of 750 million kg of raw tobacco produced annually in the country), the number of workers involved in the tobacco processing industry in India can be roughly estimated to be less than one lakh workers (Nayak, 2018). There is also a lack of information on the livelihood and health hazards of the workers in tobacco processing in India. While the hazards of habitual tobacco usage are well established, very little information is available about the effects of occupational tobacco exposure, particularly the processing of raw tobacco. The research on tobacco processing workers is a grey area. Although many attempts have been made to look into the life and working conditions of bidi rollers who are engaged in making a final tobacco product, very little information is available on processing workers who deal mainly with raw tobacco used for making tobacco products.

Studies indicate that the workers in tobacco processing industries are exposed to dust and work under unhygienic conditions without using masks and gloves and, many of them being unregistered workers without identity cards (IDs) (Mahurkar, 1990; Rudrama & Naik, 2012; Sabale et al., 2012; Khanna et al., 2013; Kaup et al., 2017; Bhalshankar & Ugle, 2020). Mahurkar (1990) reported that in many cases filed against the owners of 'Kharis' (in Gujarat processing units are known as 'Kharis'), the workers could not win as their names were not entered in the register of the unit either because they were minor or were taken on a temporary basis. It was also noted that there was no proper ventilation in the processing units. Although most of the units are registered, there was neither inspection of the working conditions nor scrutiny of the application of the labour laws in these workplaces. Bagwe & Bhisey (1993) report that although the prosperity brought by cultivation and processing of tobacco was visible by the material assets owned by the processors in the Kheda district of central Gujarat, it did not trickle down to poor workers who toiled for

hours to shower prosperity to their owners. And there are no scientific reports found off late on their improved conditions in the 'Kharis'.

A pioneering study on occupational health problems of tobacco processing workers in India carried out by Ghosh et al. (1985) revealed self-reported symptoms of vomiting, giddiness, and headache during and after processing work among 69% of workers. These workers had higher rates of nicotine and cotinine in urine excretion. Similarly, a study by Bagwe & Bhisey (1993) revealed cotinine levels in the saliva of 19% bidi workers and 100% of tobacco processing plant workers not in the habit of tobacco consumption. Mahimkar & Bhisey (1995) monitored workers engaged in the processing of bidi tobacco in India using peripheral blood lymphocytes as the test system and found increased chromosomal aberrations in tobacco processing workers compared to those in the control group, which indicates genotoxicity among tobacco processors. Bhalshankar & Ugle (2020) study found a relationship between tobacco dust exposure and changes in total thiol status. A significant decrease in levels of total thiol was found in all groups of bidi workers as compared to those in the control group. Singi & Hallikeri (2023) studied 825 tobacco processing workers in tobacco factories of Nippani city in Belagavi district of Karnataka to examine the health status of the workers. The results reveal that around 40% reported back pain caused due to being in uncomfortable posture for a prolonged time, 37% reported hand/arm fatigue 19% reported headaches, 17% reported breathing problems, 14% had vision problems and 12% reported issues with dental care. Nausea, skin diseases and palpitation were other problems reported by the workers. Despite these problems, 56% of the workers did not use protective measures like masks and gloves. In a limited study carried out by Patel et al. (2022) with fifty tobacco workers in factories of Kheda found that cough (84%), wheezing (48%), Rhinitis, back problems (40%), redness of eyes (64%), skin rashes (44%) and fever (62%) were the major health problems reported by workers.

Similar findings have emerged from studies conducted in the other countries too. The effects of occupational exposure to tobacco dust on the respiratory system of 1020 tobacco workers was attempted in Greece by Chloros et al. (2004). No significant association was found in chronic diseases in the lower respiratory systems and pollutants as against, a slightly higher reporting of disorders of upper respiratory systems in work sites compared to the participants in the control group. A study carried out by Rawan and Suzan (2023) in the city of Lattakia in Syria in a cigarette manufacturing factory revealed that more than 50% of the factory workers had respiratory problems, musculoskeletal and varicose veins problems in addition to physical and psychological fatigue and stress. However, the authors revealed that due to lack of use information on occupational hazards, there was a low reporting of occupational hazards due to environmental factors. However, it throws light on the health problems of the workers in selected factories of the district.

These studies provide useful information on the health status of tobacco workers. However, they do not explore how the conditions at the workplaces exacerbate the health hazards of the poor and unprotected workers. As we mentioned earlier, tobacco processing is largely an unorganised activity in India. It is carried out in isolated locations in closely supervised units that often accommodate the workers and their families within the premises. The current study attempts to understand the association between health problems experienced by processing workers and their conditions of living and working including access to social security benefits. It also investigates the skills these workers possess, their perceptions about the prospects of their children taking up such a hazardous activity, and the possibility of moving to other sources of livelihoods. The study aims to help policy in understanding the likelihood of future generation taking up this work, the skills required for entering alternative occupations, and the opportunities that exist for such alternatives. For the existing workers, the study identifies the gaps in availing social security benefits.

Research methodology

The study covers tobacco processing workers in Karnataka, Gujarat, Maharashtra, Andhra Pradesh (AP) and West Bengal (WB). Tobacco processing is found mainly in these five states in India.

• Sampling

The study adopted purposive sampling or snowball sampling methods chosen according to convenience in each of the selected states. Since the spread of the units and employment therein is not readily available, we choose to keep the sample to 100 tobacco processing workers in each state with a total sample of 500 tobacco processing workers. Since the number of processing workers in India is not recorded and their details are not known, we limited the sample to 100 per state. This is one of the limitations of the study. In all states actual sampling finally done was 512; 103 each in states of Andhra Pradesh, West Bengal, Gujarat, 102 in Maharashtra and 101 in Karnataka. This is because in all states 1–3 more samples have been collected by the respective teams.

It was initially decided to sample five processing units and 20 processing workers per unit selected randomly in each of the selected state for detailed interviews. However, in most of the states, we could not get entry into processing companies and the employers did not allow talking to workers. Therefore, we met workers in colonies and neighbourhoods where most of the them resided. The interviews were done at workers' homes in the morning (before going to work) and at night (after returning from work). The selected tobacco processing workers were interviewed in

Prakasam and Guntur districts of AP, Charotar region (Anand and Kheda districts) of Gujarat, Nippani tract of Belgaum district in Karnataka, Cooch Behar district in West Bengal and Sangli and Kolhapur districts in Maharashtra. These are the main centres of tobacco processing in the selected states. Given that the sample is collected through snowball method, for analysis we have used basic statistical tools and non-parametric tests for comparing groups.

The tobacco processing workers have been defined as those who are engaged in processing work for at least six months in a year and it's the major source of livelihood for the household. The tobacco processing units we visited are basically those units which process dry tobacco leaves from the farm into different grades of tobacco dust to be used in various chewing and smoking tobacco products.

Tobacco processing: Empirical findings

The tobacco processing worker was the respondent in the survey and answered most of the questions related to processing. The study covered workers of about 110 units across the five selected states – 10 in AP, 40 in Gujarat, 20 in Karnataka, 20 in Maharashtra and 20 in West Bengal. In Gujarat, tobacco processing units or *kharis* did not have names as the processing was done mainly in backyards of residences of owners of these tobacco processing units. Most of these houses and processing units are built in their farm lands which makes them ideal with required space for drying, processing and packaging the dried tobacco leaves.

• Socio-economic background and living/working conditions of the processing workers

A majority (53%) of the respondents available for discussion during the survey were female processing workers. In Gujarat, however, as high as 92% of the workers available for discussion were men. It may be noted that male migrant workers from AP constituted the largest chunk of processing workforce in Gujarat. Men formed more than half of the workers interviewed in AP and WB. In AP most of the workers were found to be intra-state migrants from the neighbouring districts. In WB, while the women mainly rolled the bidis, men do processing work.

And, 66% of the respondents in the sample were in the age group of 30 to 60 years. Around 9% of the sample processing workers were working even after the age of 60, most of them being in Karnataka and Maharashtra. In Gujarat, none of the workers were above the age of 60 possibly indicating that only able-bodied migrant workers are offered employment in the industry. Overall, 20% of the workers from the survey were widows or widowers. Nearly 81% of the sample workers were Hindus, 15% Muslims, and 4% Christians. More than half (56%) of them were illiterates with only 1% being graduates. A majority (60%) of the workers lived in

slums ranging from 17% in AP to 96% in Maharashtra. Most (73%) of the dwellings were owned by the workers, while 11% of the workers reported living in rented houses. The rest were staying in parental houses.

Liquefied Petroleum Gas (LPG) is available to 63% of households while 36% still use fuelwood as the main source and the remaining 1% use kerosene and cow dung for cooking purposes. The toilet facility is available to only 67% of the households. Others used community toilets or openly defecated.

Most of the workers belonged to weaker sections with 41% being SCs and STs. Around 37% belonged to OBCs and the remaining 28% were from the general category of the population. Around 27% were working as labour in the tobacco processing units for 10–25 years, while 17% were working for more than 25 years. While a majority started working in processing activity during the age of 25 to 50 years, around 3% had started working even after the age of 50. But it was deeply troubling to note that a few processing workers whom we interviewed, had started working as young children – even before attaining eight years of age (3%). Another 11% said that they had entered the workforce before they attained 16 years of age.


The responses of the sample processing workers on the choice of occupation reveal that processing is not a family occupation like bidi rolling, which is generally passed on inter-generationally, say, from the mother to her children, particularly, girls. Only 25% of the total sample of 512 processing workers' parents were involved in tobacco processing. Further, only in 23% of the sample households, we find other members were engaged in tobacco processing during the survey. Around 85% of the processing workers did not want their children to take up processing as their occupation (see Figure 1). And 10% of the sample processing workers reported that their children were already employed in other activities.

Only half of the processing workers are lucky to get the work throughout the year. Around 11% received work for less than three months. Around one-third of the workers in AP, WB and Maharashtra received work for more than nine months. On average around 60% of the workers work mostly 5–8 hours per day. In AP, 42% of the processing workers reported working for more than 8 hours per day.

Workers generally tie scarves or cloth on their faces to avoid dust, while a few work without it. The work is carried out from mid-October to January and from April to June. Tobacco processing is halted during monsoons to avoid dampness. As reported by the workers and the employer, the workers get a daily wage of Rs. 200 and benefits such as provident fund, pension and bonus. But these are available only to registered workers.

Most of the processing workers, except in WB, reported that their workplace had proper ventilation. Safety measures taken by processing units seem to be higher in AP as nearly half of the workers used gloves and 70% of the workers reported using

Figure 1: Distribution of Respondents by their Opinion on Children Taking Up Processing Work (%), N = 512



Source: Field Survey

masks. Usage of masks and gloves was observed to be the lowest in Gujarat. Only 18% and 31% of the workers used gloves and masks respectively while working. In 'Kharis' in Gujarat, none of the workers used gloves or masks. Most of the tasks like winnowing, grading, and packaging are carried out with bare hands in small premises surrounded by tobacco dust and residues. Nicotine observation through bare palms in the absence of gloves and exposure to dust without mask is likely to cause health problems to processing workers. It should be noted that the percentage of workers having IDs (28%), using masks (70%) and gloves (54%) is higher in AP. The working conditions seem to be better in AP as these men are involved in the processing of cigarette tobacco also. Mahurkar (1990) and Bagwe and Bhisey (1993) also indicate the harsh realities of the processing work wherein workers are denied of basic benefits and institutional recognition. Beyond these, it appears that there is no updated information regarding tobacco processing workers in the country.

Overall, only 10% of the workers possessed IDs. None of the workers in Gujarat and WB and, barely 8% of the processing workers in Maharashtra had IDs issued by the companies. This indicates the informal nature of tobacco processing wherein

workers must work at the mercy of the employer without demanding any social and economic benefits. One-fifth of the processing workers seem to be migrants. The share of migrants appears to be higher (48%) in Gujarat where it was reported that workers from Rajasthan are brought by companies specifically for processing work. A separate living arrangement is made for these workers who toil day and night within the premises of the units.


• **Sub occupations of the processing workers and Alternate employment opportunities**

Apart from processing work, workers are engaged in some other activities which are additional sources of earnings for them. The other sources of employment available to processing workers include Mahatma Gandhi National Rural Employment Guarantee Scheme (MGNREGA), reported by 3% households, agriculture labour reported by 11% households, non-agriculture labour reported by 8% households and self-employment reported by 2% households. Tobacco processing on an average gave 291 days of labour to the workers with lowest man-days of 206 in Karnataka and highest man-days of 351 days in Gujarat. This indicates that there is very little scope for processing workers to take up alternative jobs in Gujarat.

Although only 12% of the workers reported the availability of alternative employment opportunities in WB, 81% expressed their desire to shift from processing work if opportunities exist. More than half of the workers said that they were interested in shifting to alternate work in all the selected states except Karnataka (42%). In total, 56% of the workers wish to take up other works if such opportunities were available. Availability of alternate employment opportunities was reported by a higher percentage (38%) of workers in Karnataka. Overall, 30% of the workers think that alternate employment opportunities exist near their place of stay.

We tried to find the capabilities and skills of processing workers as it could help in understanding the scope for finding alternative employment opportunities available to workers in the region. Around 42% responded to the question on what employable skills they possess in addition to processing work. Of this, 56% revealed a single skill while the remaining reported possession of more than one skill. And the number of reported skills to the number of workers is in the ratio of 1:1.5 indicating that many possess additional skills. About a fourth of the workers reported being familiar with agricultural operations. This percentage corresponds with the proportion of workers who possess agricultural land (21%). Expertise in cooking was reported by 19% presumably due to the larger representation of women in the sample (53%). The other major skill reported by workers was livestock rearing (13%). As the processing workers are from rural background and agricultural families, they are aware of livestock rearing. Around 7% possess the skill of tailoring and embroidery,

Figure 2: Distribution of Workers According to the Reasons for Continuing Work in Tobacco Processing Despite Knowing that it is Harmful (%), N= 512



Source: Field Survey

2% knew bidi rolling and, less than 1% reported possessing knowledge of account keeping and computer operations, which are essential for getting employment in the current digitalized economy. Most of the business activities require the latter two skills. And, fetching sustained alternate jobs based on the skills of cooking, agriculture and livestock rearing is very difficult in rural and non-commercial areas as these are basics to many households and most of them carry out these activities for their households. They are less employable skills

• Health awareness

In response to the question how do they rate their health, only 41% of the processing workers said that they perceive their health to be good, while 45% said not so good and 14% rated their health as poor.

As reported from the field, while a majority (78%) of the processing workers knows that tobacco dust is harmful, 15% did not agree with it and 7% did not know whether it is harmful or not. But, 89.95% of the workers who felt tobacco dust was harmful across the selected five states have continued to work in the processing sector because no other work was available to them (see Fig 2). The availability of better wages is another major reason cited by 8.33% of the workers for continuing the processing work despite knowing about its adverse effects on the health of the workers.

The extant studies support the observation of high incidence of chronic illness among tobacco workers. Respiratory issues from continued inhaling of tobacco dust

and other particles, musculoskeletal disorders from prolonged awkward postures, eye/ skin irritation, headaches, and dizziness were reported in many studies. Table 1 presents the distribution of workers in the sample according to self-reported chronic illnesses. The presence of long-term illness (such as asthma, body pain, chest pain and cough, headache, acidity and gastric problems) is reported by 42% of the workers in the sample (Table 1). Self-reported illness during the reference

Table 1: Distribution of Workers According to Self-Reporting of Chronic Illness (%), N=512

States	Yes	No
West Bengal	68.93	31.07
Andhra Pradesh	80.58	19.42
Karnataka	33.63	66.37
Maharashtra	26.47	73.53
Gujarat	0.97	99.03
Total	216 (42.19)	296 (57.81)

Source: Field Survey

period of past 30 days was noted among 44% of the workers. Recent studies by Singi and Hallikeri (2023) and Patel et al. (2022) respectively in Karnataka and Gujarat also have found widespread reporting of illnesses among the processing workers.

We recognise that self-reported health status needs to be validated with bio-markers/clinical test in a larger sample for generalisation. How, it is a good indicator about feeling good physically and mentally.

We recognise that self-reported health status needs to be validated with bio-markers/clinical test in a larger sample for generalisation. How, it is a good indicator about feeling good physically and mentally.

• Wages and Social Security Benefits

Processing workers are paid in different modes – piece rates, daily wages, and monthly salaries. The wage rate for tobacco processing was found to be very low being Rs 6–7 per kg. The daily earnings varied between Rs. 150 in WB to Rs.227.49 in Karnataka (including employer's contribution towards EPF).

The owners of the processing units were worried by the labour regulations that require registered units to provide social security benefits like 10% Employee

Provident Fund (EPF), 20% bonus, holidays on national festivals, gratuity, leave with wage, and minimum wages. Hence, they do not register all the employees. They feel that it is better to close the units if all these norms are to be followed by them.

It was found that except for bonus payment, processing workers are mostly deprived of other benefits like EPF, Employee State Insurance (ESI) coverage, maternity benefits, payment of gratuity, workmen compensation in the case of death, housing facility, educational aid and group insurance. This is more so in WB where workers have not been allotted IDs. The workers in all the selected states reported receipt of bonuses though the percentage of workers receiving these benefits varies from 16% in WB to 97% in Maharashtra. Payment of bonus, however, depended on attendance. Some workers reported that their owners took them out for picnics every year at company expenses.

A few workers in AP reported the availability of many of these benefits. The payment of DA and housing facilities was reported only in Gujarat. EPF, a major social security benefit is available to only 48% of the workers in AP and 16% in Karnataka. Maternity benefit was reported by only one or two workers in AP and Gujarat. And ESI facility is reported only by 17% of the workers in AP. Around 2.5% of the workers had some kind of group insurance coverage (reported by 11% of the workers in AP and one worker in Gujarat). The payment of gratuity was reported by around 10% of the workers only in AP and Karnataka. Overall, the information available from five selected states reveals that workers in AP have most of the benefits although it is not uniform and not available to all. It could be mainly because at least 28% of the workers in AP have IDs, which means they are legally recognized.

• Annual Earnings of Tobacco Processing Workers

Tobacco processing is the main earning activity for the workers as shown in figure -3. The average annual income from tobacco processing per worker varied from a meagre amount of Rs.28,000 in Gujarat to Rs. 84,000 in AP. The earnings are higher in AP because 100% of the workers reported processing work to be available for the entire year and more than 40% reported working for more than 8 hours a day. In addition, the total sample workers in five selected states earned an average income of Rs. 15,000 annually from other activities the minimum being Rs. 7,000 in WB to a maximum of Rs. 31,000 in AP. Most (57%) of the workers take the payment weekly from their employers. Daily wagers include around 20% of the workers. Despite low wages, 27% of the workers reported some savings from their earnings.

The average income as well as median income from tobacco processing is found to be the highest in the state of Andhra Pradesh followed by West Bengal, Maharashtra, Karnataka and Gujarat in that order (Table 2). What is to be noted is the higher

share of migrant workers in Gujarat. Wages in processing are the same to both male and female in respective states and there is a great demand for processing workers as the younger generation is not inclined to processing work.

We attempted to compare the mean income of tobacco processing workers across the states if these are statistically significant or not. Kolmogorov-Smirnov (K-S) test was done to find out if the income from tobacco is normally distributed across the states and results show the statistical significance (P values) for respective K-S statistics are > 0.05 for West Bengal, Andhra Pradesh and Karnataka, indicating normal distribution but not for Maharashtra and Gujarat (< 0.05). The corresponding histograms also confirmed the same. Since we want to compare the income from tobacco processing for workers across the five states, we applied independent sample Kruskal-Wallis non-parametric test (K-W) to compare the median income for these states. It was found that the reported income from tobacco processing is statistically different across the states (N = 512, K-S statistic = 194.428, degree of freedom (df) = 4 and p-value < 0.05).

Out of 512 sample tobacco processing workers whom we interviewed, in total, 218 were males (42.6%) and rest 294 were female (57.4%). Despite women workers in our sample are largely involved in processing work and also work for longer duration in a given year than their male counterparts, we found on an average our sample male workers earn higher than female workers, Rs. 56743.6 ($\pm$ Rs. 34045.7) and Rs. 44202.9 ($\pm$ Rs. 25695.5), respectively. We applied statistical tests to determine if this difference in income from tobacco processing between male and female workers is statistically significant or not. Applying the Kolmogorov-Smirnov (K-S) test, we found the wage income from tobacco processing for our sample workers by gender was not normal.

Therefore, we have used Mann Whitney-U (MW-U) non-parametric test to compare the difference in income. It was found, the income from wage towards tobacco processing between male and female workers were different and this difference was statistically significant (N = 512, MW-U statistic = -3.310 and p-value < 0.05). All these tests were run using SPSS 16.0 and non-parametric tests were run applying auto-run option which employs appropriate model given the data. Further the data on bidi worker was collected through snowball sampling from willing processing workers in absence of a consolidated data base from which sample could have been drawn, so above non-parametric tests are apt tools to compare the group averages here.

Tobacco processing workers in addition to processing earn from their/family members' engagement as labour in MGNREGA, agriculture, bidi rolling, livestock rearing, rentals from buildings, casual labour, salaried employment and self-employment (Figure 3). The total household income of workers from several of the

Table 2: Average Annual Earnings from Tobacco Processing for Sample Workers (Rs.)

States	Mean Income ($\pm$ S.D.)	Median Income	Number of Sample Farmers
West Bengal	54894.56 $\pm$ 23324.41	54000.0	103
Andhra Pradesh	84320.39 $\pm$ 37573.30	80000.0	103
Karnataka	40769.75 $\pm$ 15423.94	40500.0	101
Maharashtra	47745.10 $\pm$ 13519.26	49400.0	102
Gujarat	28213.59 $\pm$ 18187.28	25000.0	103
Total	51236.10 $\pm$ 29888.85	48000.0	512

Source: Field Survey


reported occupations including tobacco processing was reported to be on an average Rs. 44,482 per annum and was found to be the highest in AP (Rs. 53,938) and the lowest in Gujarat (Rs. 25,254).

Since the workers are engaged in tobacco processing on an average for 291 days, there is very little scope for processing workers to take up alternative jobs. As said earlier, only those who own agricultural land (around one-fourth) have an opportunity to work in their agriculture land and rear livestock, which is complimentary to agriculture. Although women tobacco processing workers we had interviewed in these five states reported having skills of tailoring, embroidery, cooking and computer, these are not reflected in to earnings as workers are engaged in processing and household chores.

• **Are processing workers happy with their work?**

Fifty-nine percent of the processing workers reported not being happy. Of those who were not happy, a third (31%) said that they were not happy because of the sole reason that tobacco dust was harmful and they had no option to stop this activity due to absence of alternate sources of employment. Around 16% said that the wages were meagre while 11% reported health problems as the reasons for their unhappiness. Apart from these reasons, 11% reported meagre wages as well as exposure to tobacco dust as reasons for unhappiness, while 17% reported tobacco dust and health problems, 2% reported meagre wages and health problems and 12% reported all three problems as reasons for unhappiness. It should be noted that as the study reveals, around 56% of the workers (Table 3) desired to shift from tobacco processing if there exists an opportunity. Despite the non-availability of alternative

Figure 3: Different Sources of Income for Sample Tobacco Processing Workers Interviewed Across Five States (% share), N = 512



Source: Field Survey

Table 3: Tobacco Processing Workers According to their Interest to Shift to Alternative Employment (in %), N=512

States	Yes	No
West Bengal	78.64	21.36
Andhra Pradesh	55.88	44.12
Karnataka	42.00	58.00
Maharashtra	53.92	46.08
Gujarat	50.49	49.51
Total	56.19	43.81

Source: Field Survey

employment opportunities near their residence, around 79% of the workers are willing to shift from processing in WB. As against this, although more than half of the processing workers are not happy with their current work, a significant 44% are not interested in shifting. A majority being in the middle age and the non-availability of other opportunities to work near their location (the reporting of employment opportunities is lesser in states other than WB) are the reasons for their responses in this regard. Although only 30% of the workers reported availability of alternate employment opportunities near their location, 56% are willing to shift to other works if support is available. So, in the absence of alternate employment opportunities, workers may not properly answer this questions.

It should be noted that the skilling programmes of the government with respect to the provision of alternate employment to tobacco workers is negligible considering the quantum of employment in the sector, and is currently limited to bidi rollers. Both the Labour Ministry and the NGOs have targeted their skilling programmes to bidi rollers. A majority of the tobacco processing workers in our sample, do not receive social security benefits like assistance for housing, minimum wages, EPF etc., as the study found that only 10% of the workers possessed ID cards.

Summary and Policy Recommendation

There is a dearth of information on tobacco processing in India as it is largely in the unorganized sector. There is no concrete information on the location of tobacco processing units, number of units, number of workers, type of tobacco processed, internal or domestic sale, exports, etc. Processing appears to be an activity restricted to locations near the tobacco cultivation area.

The review of the available literature indicates that exposure to tobacco dust causes some adverse health effects especially respiratory outcomes such as asthma, chronic obstructive bronchitis, and allergic respiratory or nasal diseases in workers. Tobacco processing workers had self-reported symptoms of vomiting, giddiness, and headache during and after processing work and, higher rate of nicotine and cotinine in urine excretion identified in clinical studies. Many of the workers, despite being involved in hazardous work, do not receive minimum worker benefits like EPF, gratuity, bonuses, medical allowances, and minimum wages.

There appears to be a reduction in the quantum of tobacco being processed in the last 10 years. It was reported that some units have been closed, while others have shifted from three shifts to one shift and some others have become seasonal. Availability of raw materials was a major concern to tobacco processors and they assume that could be due to a reduction in tobacco production area. Electricity supply was another major issue hitting tobacco processing. More than half (58%) of the workers in the sample felt that there was a reduction in tobacco processing activity over the years. Once booming with activities, some of the processing units

now look like abandoned old castles in some of these locations visited by the survey team across five states. Workers say that anti-tobacco movement, prohibition on public smoking, restrictions on the sale and, introduction of Goods and Services Tax (GST) have cast a spell of doom on tobacco processing work. The fact that the production of tobacco remaining the same, the reported reduction in processing activity needs to be researched further for examining the real causes.

Processing is not a family occupation like bidi rolling, which is generally passed on by the mother to her children particularly the girl child. Most of the processing workers (86%) do not want their children to take up processing as their occupation. The wages for tobacco processing were found to be very low being Rs 6-7 per kg and varied between Rs. 150 in WB to Rs.227.49 per day in Karnataka. The poor working conditions, lack of social security benefits, low standard of living among a majority of workers, reporting of chronic illness among processing workers, absence of scientific studies on the health conditions of workers, and a majority of processing workers not being happy with their employment are the reasons enough for policy intervention to look into their working conditions and find out ways and means to rehabilitate them in other occupations and wean away the new entrants into tobacco processing job market.

A majority of the workers in the sample were poor slum dwellers holding BPL cards, and 20% of were widows or widowers. This highlights the vulnerable position of processing workers doomed to survive amidst the hazards of toxic dust inhalation and back-breaking hard work. Further, the fact that overall, only 10% of the workers possessed IDs indicates the informal nature of tobacco processing wherein workers must work at the mercy of the employer without demanding any social and economic benefits. Although it is a major livelihood option for processing workers who do not have access to any other alternative employment opportunity, it appears to be an exploitative work. The absence of social security benefits like housing and maternity benefits, and disparities in the availability of EPF, bonus, gratuity, etc. across the five selected states calls for bringing in uniformity in the provision of these basic social security measures to tobacco processing workers or skilling at least the younger generation who constitute around one-fourth of the processors' work force. This can enable shifting gradually from the hazardous tobacco processing.

This study should be viewed as a small step made to throw light on the living conditions and work environment of the tobacco processing in states employing a higher number of tobacco processing workers. Additional studies may be required to understand their health status in comparison to the general population using clinically diagnosed reporting and biomarkers to validate the linkage between exposure to tobacco dust and health problems among workers.

References

- Bagwe, A. N., & Bhisey, R. A. (1993). Occupational exposure to tobacco and resultant genotoxicity in bidi industry workers. *Mutation Research/Genetic Toxicology*, 299(2), 103–109.
- Bhalshankar N. & Ugle S. (2020) A research on effects of tobacco dust on status of total thiol in bidi industry workers. *Int J Adv Med*.7(8):1269–1273 [<http://www.ijmedicine.com>].
- Chloros D., Sichletidis L., Kyriazis G., Vlachogianni E., Kottakis I. & Kakoura M. (2004) Respiratory effects in workers processing dried tobacco leaves, *Allergologia et Immunopathologia*, 32(6): 344–351.
- Devi, K. R., & Naik, J. K. (2012). An Epidemiological Survey of Occupationally Exposed Beedi Workers to Tobacco Dust. *Nature, Environment and Pollution Technology*, 11(1), 135–137.
- Ghosh, S. K., Parikh, J. R., Gokani, V. N., Rao, N. M., & Doctor, P. B. (1985). Occupational health problems among tobacco processing workers: a preliminary study. *Archives of Environmental Health: An International Journal*, 40(6), 318–321.
- Kaup, S., Naseer, A., Shivalli, S., & Arunachalam, C. (2017). Occupational exposure to unburnt tobacco and potential risk of toxic optic neuropathy: a cross-sectional study among beedi rollers in selected rural areas of coastal Karnataka, India. *PLoS One*, 12(11), e0188378.
- Khanna, A., Gautam, D. S., Gokhale, M., & Jain, S. K. (2014). Tobacco dust induced genotoxicity as an occupational hazard in workers of bidi making cottage industry of central India. *Toxicology International*, 21(1), 18.
- Mahimkar, M. B., & Bhisey, R. A. (1995). Occupational exposure to bidi tobacco increases chromosomal aberrations in tobacco processors. *Mutation Research/Environmental Mutagenesis and Related Subjects*, 334(2), 139–144.
- Mahurkar U. (1990) “Tobacco workers in Gujarat labour in inhuman conditions” *The Magazine* May 15, 1990.
- Nayak, N. S. (2018). Estimates of tobacco-dependent employment in India. *Econ Polit Weekly*, 53(40), 58–62.
- Patel, J., Parmar, R., Solanki, H., Pando, B., Vohra, F., Patel, P., ... & Virendra, P. (2022). Occupational health problems among tobacco processing factory workers, at Kheda District Gujarat: A cross sectional study. *Occupational Health Problems Among Tobacco Processing Factory Workers, at Kheda District Gujarat: A Cross Sectional Study* (May 15, 2023). Jinal Patel, Rahul Parmar, Heena Solanki, Bhumi Pando, Fiza Vohra, Prachi Patel, Kailash Nagar, & Virendra Jain, 1378–1387.
- Rawan, A. S., & Suzan, Z. Assessment of Occupational Hazards Related Factors among Tobacco Workers. *Assessment*, 1(1–2023).

Sabale, R. V., Kowli, S. S., & Chowdhary, P. H. (2012). Working condition and health hazards in beedi rollers residing in the urban slums of Mumbai. *Indian Journal of Occupational and Environmental Medicine*, 16(2), 72-74.

Singi, G., & Hallikeri, A. (2023). Occupational Health status of tobacco processing workers in Nippani Taluk, Belgaum District of Karnataka: An Anthropological Study. *Antrocom: Online Journal of Anthropology*, 19(1).

Productivity and Technical Change in the Indian Pharmaceutical Sector: A Comparison of Foreign and Domestic Firms

Tulika Rohilla[#] and Boppana Nagarjuna^{*}

The effect of foreign investment on the productivity of a knowledge-intensive industry such as pharmaceuticals holds much importance. The present firm-level study uses a non-parametric mathematical linear programming model, Data Envelopment Analysis based on Malmquist Productivity Index to calculate and compare productivity, efficiency change and technical change of foreign and domestic firms of Indian pharmaceutical sector for the period 2001–2020. The findings include that foreign investment has a positive impact on the productivity of the firms. Foreign firms adapted better to the technological progress as compared to the domestic firms. The sector also experienced a regression in efficiency, especially in the years registering technological progress suggesting that a majority of firms are not able to enjoy the benefits but certain firms have progressed due to foreign investment and shifted the production frontier outwards. Domestic firms' productivity would improve through mergers and acquisitions, government support for public-private technical collaborations, expansion of research and development capabilities, and assistance in importing raw materials.

Keywords: Data Envelopment Analysis, Malmquist Productivity Index, Productivity, Technical Change, Foreign Direct Investment

[#] Corresponding Author: Tulika Rohilla, School of Economics, University of Hyderabad, Telangana, India
Email: tulikarohilla@gmail.com

^{*} University of Hyderabad, Telangana, India

The Indian pharmaceutical industry has been one of the fastest growing industries in the country since independence. According to the annual report of the Department of Pharmaceuticals (2019), it ranks tenth largest in the world in terms of value and third in volume. It accounts for 1.72 percent of the country's GDP and is projected to grow to USD 80–90 billion by 2030 (Indian Pharmaceuticals Alliance, 2019). It is one of the top FDI attracting sectors with 100 percent FDI permissible throughout automatic route in greenfield investment and 74 percent in brownfield investment.

The process of globalisation started in the 19th century and slowly engulfed the whole world. World integration is the by-product of the process. With the globalisation movement in such stride, there has been an ease in the movement of trade, flow of personnel and capital across cross-national borders. Foreign Direct Investment (FDI) has played a key role in facilitating such movements. With the remarkable development of technology, every kind of exchange, may it be business skills, production techniques or market know-how began to happen swiftly and with much ease.

The Indian pharmaceutical sector, being a high-skill, research-intensive industry has been at the centre of this transformation. At the time of independence, the Indian pharmaceutical industry did not have a prominent position. Over a period of time, it grew to be one of the most important sectors on account of a series of favourable steps taken up by the government and the initiatives of the entrepreneurs. The Patents Act, 1970 came into effect in 1972. This act provided special provisions for food, chemicals and pharmaceutical sectors which led to the abolition of product patents with the retention of process patents (Nair, 2003; Nair, 2008). It provided restrictive industrial policies which act as a stimulus in the speedy growth of the pharmaceutical sector. It also provided a base for the manufacturing of the bulk drugs as well as for the formulations (Ganguli, 2003; Nair, 2003).

The process patent regime lasted a while but since 1990s, the world has been going through a transformation regarding intellectual property protection rights in the form of Trade-Related Aspects of Intellectual Property Rights (TRIPS) Agreement. The implementation of the agreement for developing countries was done phase-wise and got completed on January 1, 2005. The objective of the transformation was to revolutionize the areas associated with intellectual property such as patents, copyrights, trademarks etc. by setting a minimum impartial level of protection for the same across the countries. TRIPS Agreement was widely accepted by the developed nations as it was seen as a tool to promote innovation and R&D in developing nations and lead to overall growth. Shapiro et al. (2014) found that stronger Intellectual Property Rights (IPRs) are associated with the transfer of the technological and innovative capabilities from developed to developing countries. This transfer generally takes place via FDI. MNCs that have patented drugs in

the developed markets are more comfortable with transferring their technology, business models and management to the countries that have stronger IPRs.

It is also a common belief that the influx of foreign investment leads to a change in firm or industry technology (Driffield & Love, 2007). The change in technical knowledge, modernization of production processes, world market integration with domestic markets and strengthening of forward and backward linkages have a positive impact on the efficiency and productivity of the firms in the sector (De Mello, 1997; Ernst, 2005). The global shift in the paradigm lead to a need for domestic firms to increase their productivity and efficiency. They took the steps of acquiring firms, new products, stepping into newer therapeutic segments, taking up better management practices and finding access to new markets (Mahajan, 2020).

Given this background, the rest of the paper has been divided as follows. Section 2 gives a brief review of earlier literature on productivity. Section 3 explains the methodology adopted in the analysis and section 4 gives the details of the data collection. Results are explained in section 5 and section 6 concludes the paper.

Literature Review

There is ample literature regarding productivity analysis which covers vast portions of theoretical as well as empirical analysis. To narrow down the literature as per the requirement of the present research, the papers highlighting the effect of foreign investment on productivity in Indian markets have been considered.

One of the most prominent researched areas while considering productivity is the productivity of the Indian manufacturing industry but the studies provide ambiguous results. Krishna & Mitra (1998) using firm-level data from different Indian manufacturing industries studied the impact of liberalization reforms on productivity and reported an increase in productivity in the time followed by the reforms. Another study by Manjappa & Mahesha (2008) examined the productivity growth and its components for manufacturing industries and found that the capital-intensive industries reported a positive productivity growth owing to positive technological change but the labour-intensive industries showed a decline in technological progress and thus a regress in productivity. Certain studies have also indicated a decline in productivity. Goldar (1986) used a multiple regression framework to understand the total factor productivity (TFP) pattern of Indian manufacturing industries in 1960–70 and recorded it be negative for majority of the firms. Another study by Balakrishnan & Pushpangadan (1994) found the productivity of Indian manufacturing firms in 1980s to be lesser than the previous decades. Jena & Chattopadhyay (2016) demonstrated positive effects of FDI on manufacturing sector provided the economy has absorptive capacity else it leads to crowding out of domestic firms.

As the Indian pharmaceutical sector is a technologically advanced sector with skilled labour and firms investing in R&D, many studies have analysed the growth, reasons and pattern of productivity of the sector. Certain studies found that the Indian pharmaceutical sector experienced technological advancements owing to technological change, investment and R&D but the benefits were not enjoyed by all firms (Mazumdar & Rajeev, 2009; Saranga & Banker, 2010; Mahajan, 2020). The implementation of TRIPS also contributed to a rise in productivity as well as profitability of Indian pharmaceutical firms (Bajaj & Nigam, 2007). Technical Efficiency Change (TEC) has been majorly found responsible for the increase in Total Factor Productivity Growth (TFPG) (Chakraborty & Pal, 2020). Firms with larger size exhibited higher productivity compared to medium and smaller enterprises (Mahajan, 2020; Das & Hoque, 2024). The level of productivity also has a direct effect on the absorption of R&D benefits of the firms (Goldar, 2013).

A certain number of studies have also highlighted the role of foreign investment on the level of productivity. As foreign firms are expected to bring better opportunities, managerial staff and technical know-how, the productivity of the firms is expected to increase. Sharma (2011) used firm level data of Indian pharmaceutical firms and stated that firms receiving FDI have better sensitivity to R&D which leads to an increase in productivity of the firms. Similar results were traced by Sharma (2015) and found out that the foreign investment received by the Indian pharmaceutical firms in the post-reform period led to a sizeable impact on the productivity of the firms. Foreign investment has also shown a positive effect on productivity growth via ownership (Thangavelu & Pattanayak, 2006), technological advancements and research efforts (Pattanayak & Thangavelu, 2011) as well as due to presence of foreign equity (Arora & Lohani, 2017). It also has a positive effect on exports which in turn leads to a rise in productivity (Mathiyazhagan & Sahoo, 2008). Foreign investment has aided the rise in productivity of the Indian pharmaceutical firms in the post-patent regime with innovation and R&D playing a decisive role (Pannu et al., 2011; Tripathy et al., 2013). Another study by Dhanora et al. (2020) examined the effect of technological innovations and foreign investment on 168 Indian pharmaceutical firms for the period 2000–2013 and found that both R&D and FDI have a positive effect on the productivity of the firms. On the other hand, an absence of horizontal spillovers from foreign firms to domestic ones has also been noted due to their ability to prevent technological outflow within the same industry (Desai et al., 2022).

As there are certain number of studies that have tried to examine the effect of foreign investment on the productivity of Indian pharmaceutical firms but most of those studies show an effect of the foreign investment on the technological innovations of the firm which in turn has an effect on the productivity of the firm. As the literature is scarce to examine the effect of foreign investment and draw a comparison with the domestic firms over a significant period of time on

the productivity of Indian pharmaceutical firms, this paper attempts to make a comparison between foreign and domestic firms of the Indian pharmaceutical sector for the period 2001–2020.

Methodology

While considering the productivity analysis, the calculation can be done either by Single Factor Productivity (SFP) or Total Factor Productivity (TFP) technique. SFP is the method of calculation by dividing the total output by only one input (Manjappa & Mahesha, 2008). It fails to provide a complete picture as it disregards the effect of other inputs in productivity of the units. It only provides partial information. TFP solves this problem as it is defined as the ratio of weighted sum of outputs to the weighted sum of inputs. It includes the effect of all the inputs on the productivity and provides comprehensive results. Total factor productivity takes into account both the changes in technical progress as well as technical efficiency.

Over a course of time, the parametric Stochastic Frontier Analysis (SFA) and the non-parametric DEA-based Malmquist Productivity Index (MPI) have become the favoured approaches for productivity analysis as they presume that the decision-making units (DMUs) are not operating at their maximum efficiency (Singh & Agarwal, 2006).

In our study, MPI has been used as it takes the efficiency of the firms into account while calculating productivity. It was first introduced by Caves et al. (1982). He used distance functions or technical efficiency functions to estimate the productivity of the firms. The technical progress and efficiency were included in the model by Fare et al. (1994). The firms in the pharmaceutical industry are expected to go through continuous technological progress which makes the adjacent period version of Fare et al. (1989) of MPI, which is defined in terms of distance functions for period t and $t+1$, the correct approach for the analysis.

The distance function measures, keeping inputs constant for period t , the maximum proportion by which outputs can be expanded for the firm in period t . Similarly, the expansion of the same output bundle relative to the frontier in the period $t+1$ is also measured. These two frontiers, for the initial as well the target period, are used to calculate the productivity changes of a firm for adjacent periods. To minimise the effect of randomly chosen technology set, the MPI production indices are calculated as the geometric mean of period t and $t+1$ ratios (Manjappa & Mahesha, 2008).

Grifell-Tatje & Lovell (1996) states certain reasons which makes MPI a popular approach in productivity analysis. The approach only uses quantity data and thus the information on input or output prices is not required. This provides a clear picture even if the information on prices is unavailable or distorted. MPI also does

not follow any particular assumption of profit maximization or cost minimization. Lastly, it decomposes the productivity indices into technical progress (outward shifting of the frontier) and the change in technical efficiency (movement along the frontier). This segregation of the indices into its two components helps in acquiring an in-depth knowledge of the whole procedure (Ma et al., 2002).

The adjacent period version of Malmquist productivity index can be expressed as

$$MI = \left[\frac{D^t(X^{t+1}, Y^{t+1})}{D^t(X^t, Y^t)} * \frac{D^{t+1}(X^{t+1}, Y^{t+1})}{D^{t+1}(X^t, Y^t)} \right]^{\frac{1}{2}}$$

Where $D^t(x^t, y^t)$ represents the distance function which gives the maximum proportion expansion of the output bundle in period t relative to the frontier in period t . Similarly, $D^{t+1}(x^t, y^t)$ is the distance function that gives the maximum proportion expansion of the output bundle in period t relative to the frontier in period $t+1$.

While measuring the productivity using adjacent periods, two separate frontiers, namely, base and final period frontiers are created. The two portions of the equation 1 represents the changes in productivity taking different periods as benchmark.

$$\frac{D^t(X^{t+1}, Y^{t+1})}{D^t(X^t, Y^t)}$$

considers the frontier for base period as the benchmark while

$$\frac{D^{t+1}(X^{t+1}, Y^{t+1})}{D^{t+1}(X^t, Y^t)}$$

captures the productivity changes keeping the frontier of the final period as the benchmark for comparison.

As there is no preference between the base and the final period frontiers as benchmarks, a geometric mean of both is taken to calculate the Malmquist Productivity Index.

Malmquist Index can be decomposed into its two components i.e. Technical Change (TC) and Efficiency Change (EC).

According to the definition of Malmquist Index:

$$MI = TC * EC$$


$$\text{Where } EC = \left[\frac{D^{t+1}(X^{t+1}, Y^{t+1})}{D^t(X^t, Y^t)} \right]$$

and,

$$TC = \left[\frac{D^t(X^{t+1}, Y^{t+1})}{D^{t+1}(X^{t+1}, Y^{t+1})} * \frac{D^t(X^t, Y^t)}{D^{t+1}(X^t, Y^t)} \right]$$

Technological progress is indicated by a value greater than one of TC and technological regress is indicated by a less than one value of TC. Similarly, EC indicates a relative shift of an inefficient firms towards or away from the frontier.

Figure 1: Decomposition of Productivity into Efficiency Change and Technical Chang



Source: Mazumdar (2013)

Mazumdar (2013) also explains the decomposition of Malmquist Index into its two components using a graph as depicted below:

Let $f(y^t)$ and $f(y^{t+1})$ be the two frontiers representing base period and final period. The position of the firm relative to base period is F and relative to final period is B.

The efficiency change of the firm is given as:

$$EC = \frac{\frac{OB}{OA}}{\frac{OF}{OE}}$$

As, while calculating technical change, MPI allows the frontier to shift in non-uniform manner, TC can be different at different inputs levels. As stated earlier, the technical change component is expressed in two portions. The first part is the TC measured relative to base period and can be expressed as

$$\frac{\frac{OB}{OC}}{\frac{OB}{OA}}$$

The TC component measured relative to the final period can be expressed as

$$\frac{OF}{OE} \bigg/ \frac{OF}{OD}$$

Therefore, the technical change component of a firm relative to MPI can be expressed as:

$$TC = \left[\frac{OF}{OE} \bigg/ \frac{OF}{OD} * \frac{OB}{OC} \bigg/ \frac{OB}{OA} \right]^{\frac{1}{2}}$$

Data

The data for the study has been collected from the Prowess package of Centre for Monitoring Indian Economy (CMIE). It is the firm- level data of Indian pharmaceutical industry for the period 2001–2020. Following Ghali & Rezgui (2001), any firm with more than 10 percent foreign equity participation is classified as a foreign firm. The study includes 38 such foreign firms. To select a comparable group of domestic firms, the asset sizes of the foreign firms were divided into ten percentile ranges. From each percentile range, domestic firms with similar asset sizes were randomly selected to ensure comparability. In total, 38 domestic firms were included in the study.

• MPI

The model used for calculating the productivity of the firms has one output and four inputs. The inputs included are labour, capital, material input and energy input. Total output is defined as the value of total sales plus change in stock which in turn is measured as opening stock minus the closing stock in output. The inputs of the model are defined as follows: (1) labour is measured in terms of wages and salaries of the workers; (2) capital is the book value of plant and machinery; (3) material input is the firm's expenditure on raw materials; and (4) energy input is the value of the firm's expenditure on power, fuel and water.

Table 1 presents the different variables and their definitions used in the model. As all the variables are expressed in monetary terms, they are appropriately deflated using price indices collected from the monthly bulletins of Reserve Bank of India (RBI). They are presented in table 2. Perpetual Inventory Method (PIM) (Balakrishnan et al., 2000) has been used to deflate capital taking 2011 as the benchmark year (Mazumdar, 2013).

Results

In the analysis of productivity, it is important to measure the extent of variation in efficiency due to technological changes. Productivity, in general terms, is defined as the ratio of outputs and inputs. It is a very similar concept to efficiency. The

Table 1: Variables and definitions used in the MPI model

Variable	Definition
Total Output	Total sales combined with change in stock (opening stock minus closing stock)
Labour	Salaries given to workers
Capital	Book value of plant and machinery
Material Input	Expenditure on raw materials
Energy Input	Expenditure in power, fuel and water

Table 2: Deflators

Variable	Deflator
Output	Wholesale Price Index for drugs and pharmaceuticals
Labour	Consumer Price Index for industrial workers
Capital	Perpetual Inventory Method
Material Input	Wholesale Price Index for chemical and chemical products
Energy Input	Wholesale Price Index for machinery and equipment

two concepts go hand –in –hand. A firm performing at a high level of efficiency would yield higher productivity. Such a firm, eventually, would cover the distance to the frontier and become one of the best performing firms in the sample under consideration. To further increase its efficiency, keeping the production frontier constant, the firm could employ more inputs to produce more outputs. This way the firm moves along the frontier. As the variable returns to scale are under consideration, the diminishing marginal returns would set in and employing more inputs no longer remains a choice. Thus, the only way to increase productivity is to experience a positive technical change and an outwards shift of the production frontier. Innovation is the prime source of such positive technical change. In conclusion, it is evident that there are two ways for a firm to experience productivity gains, one is with an increase in efficiency and the other is via positive technical change. While the first mentioned is calculated by the distance from the frontier, the latter is calculated by the shift in the production frontier.

MPI is a frontier approach that takes efficiency into consideration. It has been applied to calculate efficiency and technical change for a panel of 38 foreign and 38 domestic firms to see the effect of foreign investment on the productivity of the firms of Indian pharmaceutical sector for a period of 20 years (2001–2020).

Table 3 gives the values of the MPI and its components. A value of more than 1 for MPI suggest a percentage increment for the total factor productivity. Similarly, a value less than 1 for MPI suggest a percentage decrease in total factor productivity for an average firm. A value of 1.101 in 2004 implies that, in comparison to 2003, there was an increment of 10 percent in total productivity of foreign firms, whereas,

Table 3: Technical Change and Efficiency Change of foreign firms of Indian pharmaceutical sector, 2001–2020

Year	Total Factor Productivity	Technical Change	Efficiency Change
2001	–	–	–
2002	0.942	0.720	1.038
2003	1.024	1.085	0.944
2004	1.101	1.072	1.027
2005	1.033	1.142	0.904
2006	1.023	1.068	0.958
2007	0.857	1.014	0.845
2008	0.972	1.049	0.926
2009	1.007	0.939	1.072
2010	1.032	0.952	1.084
2011	1.137	1.056	1.077
2012	0.963	1.024	0.941
2013	1.028	1.058	0.972
2014	1.089	0.975	1.117
2015	1.010	1.079	0.936
2016	1.099	1.089	1.010
2017	1.042	1.113	0.936
2018	1.124	1.106	1.016
2019	1.040	1.083	0.961
2020	0.925	1.084	0.853

a value of 0.942 in 2002 implies that total productivity regressed by 6 percent compared to 2001. Similarly, table 4 represents the same calculations for domestic firms. The value of 1.036 for 2003 represents that total productivity increased by 3 percent in 2003 compared to 2002. A value of 0.997 in 2006 means that total productivity decreased by almost 1 percent as compared to 2005. For foreign firms, out of 19 years, total productivity shows increment for 14 years whereas for domestic firms, total productivity shows increment for 11 years. This is an initial indication that foreign investment has a positive impact on the productivity of firms. A further analysis into its components will provide a clearer picture.

Colum 3 of table 3 represents technical change for the foreign firms. A value of greater than unity represents technological progress while a value less than

Table 4: Technical Change and Efficiency Change of domestic firms of Indian pharmaceutical sector, 2001–2020

Year	Total Factor Productivity	Technical Change	Efficiency Change
2001	–	–	–
2002	0.998	0.982	1.016
2003	1.036	1.016	1.020
2004	1.032	0.958	1.078
2005	1.145	1.106	1.035
2006	0.997	0.937	1.065
2007	0.836	1.008	0.830
2008	1.226	0.985	1.244
2009	1.033	1.015	1.018
2010	1.033	1.031	1.002
2011	0.937	0.946	0.991
2012	1.040	1.059	0.982
2013	1.071	1.077	0.995
2014	1.166	0.835	1.396
2015	0.950	0.900	1.055
2016	1.005	1.046	0.961
2017	0.886	0.914	0.969
2018	1.033	1.059	0.976
2019	0.981	0.973	1.009
2020	0.905	0.939	0.964

unity represents technological regress. A value of 0.72 for technical change in 2002 implies that technology regressed by 28 percent for foreign firms as compared to 2001. A value of 1.142 for 2005 implies that relative to 2004, technology progressed or the production frontier shifted out by 14 percent. While interpreting the technical change values for domestic firms using table 4, we see a value of 0.982 in 2002 means an average firm faced a technological regression by 2 percent as compared to 2001. A value of 1.016 for 2003 suggests the production frontier shifted out by 1 percent for domestic firms as compared to 2002. Foreign firms register technological progress for 15 years and technological regress for only 4 years while domestic firms register a positive technical change for just 9 years and a regress for 10 years out of a total of 19 years.

This suggests that due to foreign investment, firms are experiencing technological progress with an outward shift in the production frontier for the majority of the years while the domestic firms are unable to adapt with technical progress and are seen facing technological regress for most of the period. Such technical progress suggests that due to investment, the sector is facing new production possibilities or new efficient firms have entered the market with superior technology.

A shift in the production frontier also increases the distance of output of firms from the frontier. This increase in distance increases average inefficiency. For foreign firms, the column 4 of table 3 represents the efficiency change component. A value of 0.944 for the efficiency change in 2003 implies that compared to 2002, the average efficiency of firms regressed by 6 percent, whereas, a value of 1.027 in 2004, implies that, on an average, the efficiency of firms has improved by 2 percent in 2004 as compared to their efficiency in 2003. The efficiency component can be interpreted in the same manner for domestic firms using table 4.

Indian pharmaceutical sector as a whole project that during the years that the industry experienced technological progress, it also witnessed a regression in its efficiency. This signifies that most of the firms are not able to enjoy the benefits that the technological progress brings along with it to the sector. As the production frontier shifts outwards, simultaneously, the fall in the level of average efficiency is also registered (Mazumdar, 2013). This implies that the technical and efficiency components in the sector have a strong negative correlation. This correlation is seen to be higher for foreign firms (79 percent) than in domestic firms (50 percent). It also suggests that a large number of firms entered the market due to lack of strong patent protection (Mazumdar, 2013). Such firms lack R&D activities and their original products. As foreign investment brings better opportunities along with it, the shift of the production frontier and also increase in inefficiency is noted more in foreign firms as compared to domestic firms.

Conclusion and Policy Recommendations

The present research extends the literature of the effect of foreign investment on the productivity and its components on the firms of the Indian pharmaceutical sector for the period 2001–2020. There are two ways for a firm to experience productivity gains, either by efficiency change (movement along the frontier) or a gain in technical change (outwards shift of the frontier). The study draws a comparison between the foreign and the domestic firms of the sector and list out the reasons for the differences in their productivities. The non-parametric MPI approach is used to decompose the productivity in efficiency change and technical change. The data comprises of 38 foreign firms and 38 domestic firms which is collected from CMIE.

The study concluded that foreign investment has a positive impact on the productivity of the firms. Foreign firms also adapted better to the technological progress as compared to the domestic firms. The sector also experienced a regression in efficiency, especially in the years registering technological progress suggesting that a majority of firms are not able to enjoy the benefits but certain firms have progressed due to foreign investment and shifted the production frontier outwards. Thus, we conclude that foreign investment is paramount for firms to perform better and improve their productivity but certain policy changes are required for the whole sector to enjoy the benefits brought along with it. The firms would benefit from mergers and acquisitions as it would overall increase their size which in turn would increase their absorptive capacity and lead to reap better benefits from investments. With an increase in size, the firms might also be able to use the unutilised plant and machinery and this might lead to capital efficiency as well.

Smaller firms do not possess the resources necessary to engage in importing which would also lead to increase in productivity. These firms might receive assistance from the government in one of two ways: either by providing financial assistance or by establishing markets. These markets would facilitate firms to do business by purchasing medicines at a lower price which would help the firms in recouping the money that they invested.

Research and development is also known to be an important determinant in productivity enhancement. Hence, it is recommended that all businesses invest in R&D. This investment should be done in order to develop new goods as well as processes. Scale economies are also present for firms that invest in R&D. However, not all firms are able to absorb foreign investment by assuming the risk of investing in R&D-related activities. In order to facilitate these processes, the government might engage in public-private technical collaborations and contribute to the expansion of research and development endowment, which in turn would assist in more effective utilisation of the received foreign investment.

References

- Arora, N., & Lohani, P. (2017). Does foreign direct investment spillover total factor productivity growth? A study of Indian drugs and pharmaceutical industry. *Benchmarking an International Journal*, 24(7), 1937–1955.
- Bajaj, A. K., & Nigam, S. (2007). Globalization in the Indian Pharmaceutical Industry—FDI spillovers and implications on Domestic Productivity: 1991–2007. is a research project done under IIM Ahmedabad.
- Balakrishnan, P., & Pushpangadan, K. (1994). Total factor productivity growth in Indian manufacturing: a fresh look.

Balakrishnan, P., Pushpangadan, K., & Babu, M. S. (2000). Trade liberalisation and productivity growth in manufacturing: Evidence from firm-level panel data. *Economic and Political weekly*, 3679–3682.

Caves, D. W., Christensen, L. R., & Diewert, W. E. (1982). The economic theory of index numbers and the measurement of input, output, and productivity. *Econometrica*, 50(6), 1393.

Chakraborty, C., & Pal, D. (2019). Total Factor Productivity Growth of Indian Pharmaceutical Industry: A Non-Parametric Approach. *Vidyasagar University Journal of Economics*, XXIV, 129–131.

Das, S., Hoque, A. Firm-level productivity and its determinants in the Indian pharmaceutical industry. *Decision* 50, 439–459 (2023).

De Mello, L. R. (1997). Foreign direct investment in developing countries and growth: A selective survey. *The Journal of Development Studies*, 34(1), 1–34.

Desai, G., 1, Srinivasan, P., 2, & Gowda, A. B., 3. (2022). FDI spillover effects on the productivity of the Indian pharmaceutical industry: Panel data evidence. *Journal of Asian Finance, Economics and Business*, 9(8), 0109–0121.

DH, M. (2008). Productivity Performance of Selected Capital-Intensive and Labor-Intensive Industries in India During Reform Period: An Empirical Analysis. *ICFAI Journal of Industrial Economics*, 5(4).

Dhanora, M., Danish, M. S., & Sharma, R. (2021). Technological innovations and firms' productivity in new patent regime: Evidences from Indian pharmaceutical industry. *Journal of Public Affairs*, 21(1), e2136.

Driffield, N., & Love, J. H. (2007). Linking FDI motivation and host economy productivity effects: conceptual and empirical analysis. *Journal of International Business Studies*, 38(3), 460–473.

Ernst, C. (2005). The FDI – employment link in a globalizing world: The case of Argentina, Brazil and Mexico. *Employment Strategy Papers*. https://www.ilo.org/wcmsp5/groups/public/---ed_emp/---emp_elm/documents/publication/wcms_114029.pdf

Faere, R., Grosskopf, S., Lovell, C. a. K., & Pasurka, C. (1989). Multilateral Productivity Comparisons When Some Outputs are Undesirable: A Nonparametric Approach. *The Review of Economics and Statistics*, 71(1), 90.

Färe, R., Grosskopf, S., Norris, M., & Zhang, Z. (1994). Productivity Growth, Technical Progress, and Efficiency Change in Industrialized Countries. *The American Economic Review*, 84(1), 66–83. <http://www.jstor.org/stable/2117971>

Ganguli, P. (2004). Patents and patent information in 1979 and 2004: a perspective from India. *World Patent Information* 26, 61–62.

Ghali, S., & Rezgui, S. (2011). FDI Contribution to Technical Efficiency in the Tunisian Manufacturing Sector: Evidence from Micro-panel Data. *International Economic Journal*, 25(2), 319–339.

Goldar, B. (1986). Import Substitution, Industrial Concentration And Productivity Growth In Indian Manufacturing. *Oxford Bulletin Of Economics And Statistics*, 48, 2.

Goldar, B. (2013). R&D intensity and exports: a study of Indian pharmaceutical firms. *Innovation and Development*, 3:2, 151-167.

Government of India, Ministry of Chemicals & Fertilizers, & Department of Pharmaceuticals. (2019). Annual Report 2019-20.

Grifell-Tatjé, E., & Lovell, C. (1996). Deregulation and productivity decline: the case of Spanish savings banks. *European Economic Review*, 40(6), 1281-1303.

Indian Pharmaceutical Alliance. (2019). The Indian pharmaceutical industry – the way forward.

Jena, P. K., Chattopadhyay, U., School of Economics, Shri Mata Vaishno Devi University, Jammu & Kashmir, India, & National Institute of Industrial Engineering (NITIE), Mumbai. (2016). Impact of Foreign Direct Investment (FDI) Inflows on Productivity: Evidence from Panel Data Analysis. In *Productivity [Journal-article]*.

Krishna, P., & Mitra, D. (1998). Trade liberalization, market discipline and productivity growth: new evidence from India. *Journal of development Economics*, 56(2), 447-462.

Ma, J., Evans, D. G., Fuller, R. J., Stewart, D. F., International Technologies Centre, & Faculty of Architecture, Building & Planning, The University of Melbourne. (2002). Technical efficiency and productivity change of China's iron and steel industry. In *Int. J. Production Economics* (Vol. 76, pp. 293-312).

Mahajan, V. (2019). Is productivity of Indian pharmaceutical industry affected with the introduction of product patent act? *Indian Growth and Development Review*, 13(1), 227-258.

Mathiyazhagan, M. K., & Sahoo, D. (2008). Do foreign direct investment inflows benefit the major sectors in India. *Institute of South Asian Studies*.

Mazumdar, M. (2013). Performance of pharmaceutical companies in India [English]. *Physica-Verlag*.

Mazumdar, M., & Rajeev, M. (2009). A comparative analysis of efficiency and productivity of the Indian pharmaceutical firms: a Malmquist-meta-frontier approach. *Bangalore, India*.

Nair, G. (2003). Indian patent law and pharma industry.

Nair, G. N., Ph. D. & Gopakumar Associates. (2008). Impact of TRIPS on Indian pharmaceutical industry. In *Journal of Intellectual Property Rights: Vol. Vol 13* (pp. 432-441).

Pannu, H., Kumar, U. D., & Farooque, J. A. (2011). Efficiency and productivity analysis of Indian pharmaceutical industry using data envelopment analysis. *International Journal of Operational Research*, 10(1), 121

Samirana Pattnayak, S. and Thangavelu, S.M. (2011), "Linkages and technology spillovers in the presence of foreign firms: Evidence from the Indian pharmaceutical industry", *Journal of Economic Studies*, Vol. 38 No. 3, pp. 275–286.

Saranga, H., & Banker, R. D. (2010). Productivity and technical changes in the Indian pharmaceutical industry. *Journal of the Operational Research Society*, 61(12), 1777–1788.

Shapiro, R. J., & Mathur, A. (2014). How India can attract more foreign direct investment, create jobs and increase GDP: The benefits of respecting the intellectual property rights of foreign pharmaceutical producers. *SSRN Electronic Journal*.

Sharma, C. (2011). RD and productivity in the Indian pharmaceutical firms. Munich Personal RePEc Archive.

Sharma, C. (2016). R&D, Technology Transfer and Productivity in the Indian Pharmaceutical Industry. *International Journal of Innovation Management* Vol. 20, No. 1.

Singh, S. P., & Agarwal, S. (2006). Total factor productivity growth, technical progress and efficiency change in sugar industry of Uttar Pradesh. *The Indian Economic Journal*, 54(2), 59–82.

Tripathy, I. G., Yadav, S. S., & Sharma, S. (2013). Efficiency and productivity in the process and product patent regimes: empirical evidence from the Indian pharmaceutical industry. *International Journal of Economics and Business Research*, 6(1), 1.

Thangavelu, S. M., & Pattnayak, S. S. (2005). Linkages spillovers and foreign ownership: evidence from the Indian pharmaceutical firms. In 32nd European Association of Research in Industrial Economics (EARIE) Conference.

Burden of Distress Financing for Hospitalization in India: Prevalence and Patterns from Household Health Care Consumption Survey, 2017–18

Sunil Rajpal[#], Sneha Gupta^{*}, Shreya Ronanki^{*}

The Indian health system is predominantly characterized by out-of-pocket healthcare expenditure and limited insurance coverage. While affluent households typically finance treatment through income and savings, economically disadvantaged households to rely on distressed sources like selling assets and borrowing to meet the healthcare needs. This study examines the levels and patterns of distress financing for hospitalization care in India across various demographic and socioeconomic groups. Furthermore, it compares the prevalence of distress financing across major ailment categories. The analysis utilizes data from the 75th round of the National Sample Survey 2017–18 to examine the overall prevalence of distress financing and the final analytic sample consists of 66237 participants. The key respondents include individuals who were hospitalized within the last 365 days. The binary outcome variable indicates whether patients funded their inpatient care through household income/savings (o), or distress means borrowings, sale of physical assets, contributions from friends and relatives, and other sources. The explanatory variables include MPCE quintiles, education, and place of residence. Four ailment categories were cancer, cardiovascular diseases, diabetes, and accidents. The analysis indicates that overall, 43.9% of inpatient cases relied on distress means of financing as either primary or secondary source. Across private and public facilities, the prevalence of distress financing was 35.9% and 49.4% respectively. Within private hospitalization facilities, a sharp gradient was observed across MPCE quintiles with a prevalence of 58.5% amongst the poorest as compared to 35.0% amongst the richest households. These findings highlight the need for strengthened social protection policies and expanded healthcare coverage to alleviate the reliance on distressed healthcare financing. Furthermore, prioritizing insurance-based healthcare systems is critical to achieving equitable and sustainable reductions in the financial burden of healthcare.

[#] Corresponding Author: Sunil Rajpal, FLAME University, Pune, India
Email: sunil.rajpall@flame.edu.in

^{*} FLAME University, Pune, India

Rising healthcare costs impose a significant financial burden, particularly on low-income households, who are disproportionately vulnerable. In developing countries, over one-third of healthcare expenditure is financed through out-of-pocket (OOP) expenditure by households. The rising burden of out-of-pocket health expenditure is one of the prominent health policy concerns, particularly in low-income and densely populated countries like India. Universal Health Coverage is, therefore, is one of the primary health agendas to achieve two fundamental objectives: (a) equity in healthcare access, and (b) financial protection from the high treatment cost of non-communicable diseases (NCDs). Additionally, studies before the implementation of the National Health Mission (NHM) have shown a distinct pro-rich bias towards access and utilization of healthcare facilities in India. Moreover, the OOP healthcare expenditure for those seeking care in private facilities has increased exponentially in India, thereby worsening the economic challenges faced by financially disadvantaged households. It is important to highlight that the treatment cost of chronic diseases like cancer and cardiovascular diseases (CVDs) is substantially higher, often driving households into economic hardship and compelling them to distress modes of financing. Distressed financing encompasses borrowings, contributions (with or without repayment obligations), and the sale of assets by households to address healthcare expenses. Previous studies indicate that approximately 60% and 32% households in India rely on borrowings and contributions from friends and relatives, respectively, for financing cancer hospitalization in India (Joe, 2015). The widespread reliance on distress modes of financing reflects the regressive nature of the healthcare system and calls for urgent policy intervention. Similar findings from other low-income countries corroborate that unpredictable and high OOP payments on healthcare lead to severe financial distress among households (Bonu et al., 2005; Bonu et al., 2009; Narayan et al., 2000a; Narayan et al., 2000b; Sauerborn et al., 1996; Damme et al., 2004).

In response to the critical need to address these challenges, the Government of India has formulated policies emphasising financial protection and equitable access to healthcare services. The National Health Policy (NHP), 2017, accorded 'affordability' as a key principle. For instance, the NHP states that "[a]s costs of care increase, affordability, as distinct from equity, requires emphasis. Catastrophic household health care expenditures, defined as health expenditure exceeding 10% of its total monthly consumption expenditure or 40% of its monthly non-food consumption expenditure, are unacceptable" (MoHFW, 2017). One crucial issue requiring immediate policy attention is the financing of healthcare, particularly for the elderly population, who are disproportionately exposed to chronic non-communicable diseases (NCDs). The rise in healthcare expenditures is primarily driven by the combined effects of an increasing aging population and the growing prevalence of NCDs. The age factor in the increasing prevalence of NCDs is inevitable, as adults and the elderly are more susceptible to chronic diseases. Moreover, the

treatment of NCDs, particularly cardiovascular diseases (CVDs), is not only prolonged in duration, but also incurs high costs, with comparatively lower survival rates. In addition to the fact that India, like many other developing countries, lacks infrastructure to accommodate the increasing burden, policymakers must prioritize addressing the issue of skyrocketing health spending by households, financed through distress modes of financing at the household level.

There is a dearth of studies examining the incidence of distress financing for healthcare for both public and private health facilities. Although previous studies have underscored the significant burden of OOP expenditure, there is a notable lack of evidence investigating distress financing patterns independently for public and private health care facilities in India, particularly on a nationwide basis. In addition to this, given the wide array of improvements in health system financing and infrastructure, it is critical to examine the patterns in distress modes of financing healthcare separately for public and private hospitals. Although the NHP 2017 explicitly prioritizes healthcare affordability, it is essential to assess the role of both public and private health facilities in the overall burden of healthcare-related borrowings in India. Furthermore, it will also help determine whether investments in public health care have significant protective effects on household health financing and whether these interventions need to be further strengthened.

Drawing on the data from the health round of the National Sample Survey (2017–18), this paper aims to examine the incidence and modes of hospitalization financing at the household level. Beyond examining prevalence, the study explores how key socioeconomic correlates are associated with distressed healthcare financing (borrowings, sale of assets, and contributions from friends or relatives). The analysis uncovers the complex dynamics of distress modes of financing across the intersections of multiple factors, including gender, social categories, and economic status, that can have a significant impact while deciding upon the use of various sources of healthcare financing (Van Doorslaer et al., 2007). In this regard, a small area study on Koppal district in Karnataka in India provides compelling evidence that economically challenged men had better access to credit markets and were more likely to take loans or sell assets to finance healthcare expenses (Sen & Iyer, 2012). Even though the prevailing evidence comes from small-area studies, from a policy perspective, it is essential to analyse the broader scope, proportionate impact, and correlates of distressed healthcare financing.

Data and Methods

• Data

This study draws upon nationally representative data from Social Consumption: Health survey (75th round) of India (NSSO, 2018). The survey was conducted in 2017–18 by the National Sample Survey Organization (NSSO), Ministry of Statistics

and Program Implementation, Government of India. The Social Consumption: Health survey collects information on morbidity, treatment-seeking, and financing of hospitalization (inpatient) and ambulatory (outpatient) care services for reference periods of 365 days and 15 days, respectively. The survey also details ailments treated through medical care, the extent of utilization of Government hospitals, and treatment-related expenditures in public and private sectors. Additionally, the survey provides household-level characteristics on demographics and access to services and utilities as well as individual-level data on age, sex, education, monthly per capita expenditure, and primary occupation.

- **Survey Design**

The Social Consumption: Health Survey covers a representative sample of households adopting a stratified multi-stage survey design covering India. A rural/urban stratification is created within clusters called state regions, which constitute a contiguous group of districts within a State or union territory with similar features. Within each district of a State/Union Territory, rural and urban strata are formed. The first-stage units are selected using the circular systematic sampling system of census-identified villages (rural sector) and urban frame survey blocks (urban sector) of each district. Villages and blocks with larger sample sizes are divided into multiple “hamlet groups” or “sub-blocks” and households belonging to only two “hamlet groups” and one randomly selected “sub-blocks” are selected as part of the second stage sampling process. The households within each village are further categorized into two strata based on their level of affluence and are then circular systematically selected to form the final sample. The 75th round of the Morbidity and Healthcare Survey covers a sample of 113822 households and 555351 individuals.

- **Outcomes**

We investigated the percentage of individuals largely relying on distress financing mechanisms to receive inpatient treatment. The survey collects information on the major source of financing to capture whether the majority of out-of-pocket expenditure was incurred through distressed means or not. Components such as borrowings (with or without interest), contribution from friends and relatives (with or without repaying option), and sale of assets are together defined as distressed financing. We further categorize it into “First source” and “Second source” to highlight the differences in cases that rely on distress mechanisms as the primary mode of financing and those that do so as a supplementary source. “Either source” reflects the total burden of distress financing.

- **Socioeconomic Indicators**

The study focuses on indicators of socioeconomic status including household monthly per capita expenditure (MPCE) quintile, education, and social group of

the patient. Years of education was used to categorize the patients as illiterate (no formal schooling), upto primary education (1–5 years), upto middle school education (6–10 years), secondary education (11–12 years) and higher education (graduate school and above). Social group was categorized into the three historically marginalized groups of scheduled tribes (ST), scheduled castes (SC), other backward classes (OBC), alongside other castes. In addition, we also include information on place of residence (urban vs. rural), sex of the patient, religion (Hinduism, Muslim, Christianity, others), and region.

Statistical Analyses

The study reports the of prevalence of distress financing for inpatient care across socioeconomic categories. The concentration index (CI) is used to discern the socioeconomic gradient in inpatient care (Wagstaff et al., 1991; Erreygers, 2009) with a focus on public and private hospitals separately. The value of CI ranges between +1 and –1 with zero depicting no inequality and large positive values indicating greater concentration of distress financing cases among the richer households. Further, we employ logistic regression (adjusting for state and community-level fixed effects) to understand the mutually adjusted associations of distress financing prevalence with various socioeconomic indicators in a multivariate framework. The logistic regression estimates are reported in the form of Odds Ratio (OR) with 95 percent confidence intervals. These OR are the relative measure of effect which allows comparisons of groups relative to the reference group. The analysis was carried out in Stata 15 (StataCorp, 2013; Leckie & Charlton, 2013). All the analyses use multipliers as prescribed by the NSSO (NSSO, 2018).

Results

About 45.7% of inpatient cases were reported to resort to distress modes of financing with 18.9% and 29.2% as a primary and secondary source respectively (Table 1). Compared to the highest MPCE quintile, resorting to distress financing was more common among poorer sections. For instance, 37% of inpatient cases in the fifth quintile against 49.5% in the first quintile went for distress financing sources.

Across public and private facilities, 37.8% and 51.2% of hospitalization cases were financed through distress sources respectively (Table 2). For private hospitals, a clear gradient was observed across MPCE quintiles with 60.6% prevalence among the poorest households and 37.6% among the richest households. However, such a gradient was not observed in the case of public hospitals. We could not find notable differences in the prevalence across genders in both public and private

facilities. For private hospitalization cases, socially affluent sections are less prone to finance through distress modes compared to categories with social reservations. For example, the prevalence among cases from ST households and unreserved households was 56.9% and 43.2% respectively. Such a pattern was missing in the case of inpatient cases in public hospitals. Interestingly, private inpatient cases also depict a stark difference across rural (57.2%) and urban (42.0%) areas in the prevalence of distress financing.

Across states, Andhra Pradesh reported to have the highest percentage (64.2%) of inpatient cases resorting to distress sources of financing followed by Manipur (62.3%), Goa (59.5%), Arunachal Pradesh (59.4%), and Karnataka (59.3%) (Fig1 – All). On the other hand, states like Himachal Pradesh (23.5%), Gujarat (30.3%) and Madhya Pradesh (31.1%) had relatively much lower burden. These findings were consistent across public and private facilities with higher magnitude in private hospitalization cases. For instance, the north-eastern states of Nagaland (Public: 63.9%; Private: 82.6%), and Manipur (Public: 59.8%; Private: 72.2%) had a much higher burden of distress financing.

We also quantified the magnitude of distress financing burden across major ailment groups for inpatient treatment, i.e., CVDs, Diabetes, Cancer, and Accidents. The prevalence of distress financing was highest for Cancer (67.8%), followed by accidents (52.6%), diabetes (47.4%), and CVDs (47.4%) (Fig 2). This pattern was consistent for both the first source and the second source. For private hospitals, the magnitude of burden was relatively higher for all groups with 72.9% of cases with distress financing for cancer, 61.2% for accidents, 53.4% for CVDs, and 50.7% for diabetes. In the case of public hospitals as well, distress financing was highest for cancer inpatient cases, followed by accidents and CVDs.

Estimates from the concentration index depict a pro-rich bias with a higher burden of distress financing for inpatient cases among poor households (Fig3). Econometric estimates from logistic regression reflect higher odds of distress financing for the treatment of adults (OR: 1.30; 95% CI: 1.24; 1.36) and the elderly population (OR: 1.24; 95% CI: 1.15; 1.31) compared to the younger cohort (Table 3). Across the MPCE quintiles, the likelihood of relying on borrowings and other distress means for adult hospitalization was significantly higher among households in the poorest (OR: 1.53; 95% CI: 1.44; 1.63) and poorer households (OR: 1.48; 95% CI: 1.39; 1.57) compared to those in the richest quintile.

These observations were consistent for both public and private facilities. For inpatient treatment of cancer, distress financing was reported as two times more likely than those with any other ailments (OR: 2.71; 95% CI: 2.36; 3.12). The probability of distress financing for inpatient treatment was reported significantly higher for rural areas (OR: 1.24; 95% CI: 1.19; 1.29) compared to urban areas. We also assessed the econometric association of distressed financing cases with the

Table 1. Prevalence (%) of Distress Modes of Financing Hospitalization Care by Socioeconomic Background, India, NSS 2017-18

Characteristics	First Source			Second Source			Either Source		
	Proportion (%)	Standard Error	Sample (N)	Proportion (%)	Standard Error	Sample (N)	Proportion (%)	Standard Error	Sample (N)
Age group (years)									
0-15	16.75	1.01	11207	25.96	0.50	11207	41.22	0.65	11207
15-60	19.24	0.43	41170	29.75	0.28	41170	46.60	0.33	41170
60+	19.93	0.97	13860	29.82	0.58	13860	46.74	0.68	13860
MPCE Quintiles									
Poorest	18.59	0.80	9030	33.54	0.59	9030	49.46	0.66	9030
Poorer	19.59	0.76	11139	31.94	0.54	11139	49.04	0.63	11139
Middle	21.58	0.79	13070	31.06	0.50	13070	49.46	0.60	13070
Richer	19.68	0.82	16415	28.09	0.48	16415	45.61	0.58	16415
Richest	15.48	1.07	16583	23.08	0.47	16583	37.04	0.55	16583
Education									
Illiterate	21.56	0.61	18864	29.38	0.41	18864	48.14	0.49	18864
Primary	14.35	3.36	463	33.44	2.73	463	45.68	3.01	463
Secondary	17.82	3.81	351	25.75	3.29	351	42.27	3.61	351
Higher	17.86	0.48	46559	29.03	0.28	46559	44.69	0.33	46559
Gender									
Male	20.25	0.57	34589	29.57	0.33	34589	46.95	0.39	34589
Female	17.64	0.48	31642	28.70	0.32	31642	44.49	0.37	31642
Social Group									
Scheduled Tribes	16.74	0.71	7425	30.73	0.62	7425	44.34	0.69	7425
Scheduled Castes	20.77	0.89	11090	31.35	0.60	11090	49.15	0.68	11090
Other Backward Classes	20.30	0.64	26641	30.52	0.37	26641	48.38	0.43	26641
Others	16.51	0.70	21081	25.63	0.38	21081	40.35	0.47	21081
Religion									
Hinduism	18.85	0.43	50204	29.24	0.26	50204	45.81	0.31	50204
Islam	19.92	1.09	8997	28.89	0.62	8997	46.17	0.78	8997
Others	18.58	1.08	7036	28.55	0.74	7036	44.17	0.86	7036
Sector									
Rural	20.54	0.48	36862	31.27	0.32	36862	49.00	0.37	36862
Urban	16.12	0.57	29375	25.25	0.32	29375	39.81	0.38	29375
All	18.98	0.37	66237	29.15	0.23	66237	45.76	0.27	66237

Table 2. Prevalence (%) of Distress Modes of Financing (First or Second) by Public and Private Hospital Facilities across Socioeconomic Background, India, NSS 2017-18

Characteristics	Public			Private		
	Proportion (%)	Standard Error	Sample (N)	Proportion (%)	Standard Error	Sample (N)
Age group (years)						
0-16	33.14	1.02	5432	46.81	0.80	5775
15-60	38.25	0.43	19222	52.52	0.43	21948
60+	40.44	0.98	5935	50.91	0.85	7925
MPCE Quintiles						
Poorest	38.98	0.80	5221	60.64	0.95	3809
Poorer	38.63	0.76	6115	59.26	0.92	5024
Middle	38.87	0.80	6586	57.67	0.79	6484
Richer	36.56	0.83	7569	51.83	0.75	8846
Richest	35.75	1.07	5098	37.36	0.61	11485
Education						
Illiterate	38.59	0.61	9399	55.87	0.67	9465
Primary	35.11	3.52	280	57.95	4.81	183
Secondary	35.59	3.81	194	48.79	6.01	157
Higher	37.57	0.48	20716	49.24	0.41	25843
Gender						
Male	39.30	0.58	15608	52.27	0.48	18981
Female	36.44	0.48	14977	50.06	0.51	16665
Social Group						
Scheduled Tribes	37.51	0.72	5369	56.86	1.43	2056
Scheduled Castes	40.21	0.90	5951	58.06	0.89	5139
Other Backward Classes	38.57	0.65	10855	54.45	0.53	15786
Others	35.01	0.71	8414	43.16	0.58	12667
Religion						
Hinduism	37.41	0.43	21938	51.61	0.39	28266
Islam	38.57	1.10	4743	52.63	1.01	4254
Others	43.21	1.09	3908	44.57	1.29	3128
Sector						
Rural	38.83	0.48	19225	57.20	0.51	17637
Urban	35.63	0.58	11364	42.01	0.47	18011
All	37.88	0.38	30589	51.22	0.35	35648

intersection of social category and MPCE quintiles (Fig 4). A clear gradient was observed with a significantly higher likelihood of resorting to distress means among the socially deprived and poorest MPCE quintile households in both public and private hospitals (Fig 4). Further, compared to the patients with the highest education from the richest households, the odds for patients from less educated backgrounds from poorer households have a significantly higher probability of resorting to borrowings and other (Fig 5).

Figure 1: Prevalence of Distress Modes of Financing for Hospitalization Care by Type of Facilities and States, India, NSS, 2017-18



Figure 2: Prevalence (%) of Distress Modes of Financing Hospitalization Care by Broad Ailment Categories, India, NSS 2017-18




Figure 3: Prevalence (%) of Distress Modes of Financing by Public and Private Hospital Facilities across Rural and Urban sectors, India, NSS 2017-18




Table 3. Econometric association between distress financing for hospitalization and socio-economic correlates, by public and private hospitals, India, NSS 2017-18

Correlates	Public		Private		All	
	Odds Ratio	95% CI	Odds Ratio	95% CI	Odds Ratio	95% CI
Age group (years)						
0-15 ®	-	-	-	-	-	-
15-60	1.24***	[1.16 ; 1.33]	1.39***	[1.30 ; 1.48]	1.30***	[1.24 ; 1.36]
60+	1.28***	[1.07 ; 1.40]	1.26***	[1.16 ; 1.36]	1.24***	[1.16 ; 1.31]
Gender						
Female ®	-	-	-	-	-	-
Male	1.13***	[1.07 ; 1.87]	1.15***	[1.10 ; 1.21]	1.15***	[1.11 ; 1.19]
Sector						
Urban ®	-	-	-	-	-	-
Rural	1.17***	[1.10 ; 1.24]	1.36***	[1.29 ; 1.43]	1.24***	[1.19 ; 1.29]
MPCE Quintiles						
Richest ®	-	-	-	-	-	-
Poorest	1.63***	[1.48 ; 1.80]	1.78***	[1.62 ; 1.95]	1.53***	[1.44 ; 1.63]
Poorer	1.53***	[1.39 ; 1.68]	1.69***	[1.56 ; 1.83]	1.48***	[1.39 ; 1.57]
Middle	1.41***	[1.29 ; 1.54]	1.61***	[1.50 ; 1.74]	1.43***	[1.35 ; 1.51]
Richer	1.20***	[1.10 ; 1.30]	1.38***	[1.29 ; 1.47]	1.25***	[1.19 ; 1.31]
Education						
Higher ®	-	-	-	-	-	-
Illiterate	1.03	[0.97 ; 1.09]	1.23***	[1.16 ; 1.30]	1.12***	[1.07 ; 1.16]
Primary	1.25	[0.97 ; 1.61]	1.29	[0.95 ; 1.76]	1.18	[0.97 ; 1.43]
Secondary	1.29	[0.95 ; 1.74]	1.14	[0.80 ; 1.61]	1.13	[0.90 ; 1.42]
Social Group						
Others ®	-	-	-	-	-	-
Scheduled Tribes	1.12	[1.02 ; 1.22]	1.51***	[1.36 ; 1.68]	1.14***	[1.07 ; 1.22]
Scheduled Castes	1.17***	[1.08 ; 1.27]	1.12	[0.99 ; 1.25]	1.36***	[1.29 ; 1.43]
Other Backward Classes	1.19***	[1.12 ; 1.28]	0.96	[0.87 ; 1.07]	1.40***	[1.34 ; 1.46]
Diseases						
Cancer- No ®	-	-	-	-	-	-
Cancer- Yes	3.03***	[2.45 ; 3.75]	2.45***	[2.04 ; 2.95]	2.71***	[2.36 ; 3.12]
Cardiovascular Diseases- No ®	-	-	-	-	-	-
Cardiovascular Diseases- Yes	1.16***	[1.05 ; 1.27]	1.22***	[1.13 ; 1.32]	1.20***	[1.12 ; 1.27]
Diabetes- No ®	-	-	-	-	-	-
Diabetes- Yes	1.17	[0.89 ; 1.29]	0.93	[0.80 ; 1.09]	1.00	[0.89 ; 1.13]
Accidents- No ®	-	-	-	-	-	-
Accidents- Yes	1.07***	[1.08 ; 1.26]	1.47***	[1.37 ; 1.58]	1.34***	[1.28 ; 1.41]
N	27,526		31,732		58,209	

Note: Odds ratios are estimated employing logistic regression, adjusting for insurance status, chronic ailment (any), religion, and state-fixed effects. Estimations are *significant at 0.10, ** at 0.05, *** at 0.01 level. ® refers to the reference category for the correlates.

Figure 4: Econometric association between distress financing for hospitalization and Social category*MPCE Quintiles, by public and private hospitals, India, NSS 2017-18





Figure 5: Econometric association between distress financing for hospitalization and Education Groups*MPCE Quintiles, by public and private hospitals, India, NSS 2017-18



Discussion

The study identifies four key findings. First, nearly half of the inpatient cases in India rely on distress financing mechanisms to meet healthcare expenses, with a disproportionately higher burden in rural areas. Second, the reliance on distress financing is substantially more pronounced for treatments sought in private hospitals compared to public ones, a trend that remains consistent across various demographic and socioeconomic strata. Third, the econometric analysis reveals a significantly higher probability of resorting to distress financing for inpatient treatment among socioeconomically vulnerable households, indicating borrowing as a potential marker of economic vulnerability. Fourth, the burden for distress financing was particularly higher for hospitalization related to cancer treatments across the broad ailment categories. Notably, inpatient treatment for accident cases also exhibited a substantial reliance on distress financing. This could be attributed to the fact that cancer treatment often entails prolonged and expensive medical care, which burdens households with limited financial resources, thereby rising their dependence on distress financing. Whereas, accident injuries necessitate immediate and unplanned medical interventions, leading to sudden financial pressures that force households to resort to distress financing mechanisms.

Our findings are consistent with previous studies on the subject matter. Existing literature from low-income countries highlights that a notable proportion of households incur financial debt or sell household assets to manage healthcare expenses (Sauerborn et al., 1996; Wagstaff et al., 1991; Leive & Xu, 2008; Kruk et al., 2009). For instance, the World Health Survey (2002–04) conducted across several low- and middle-income countries, reported a mean prevalence of borrowing and asset sales at 22% and 10%, respectively (Kruk et al., 2009). In light of these figures, it is expected that households in a low-income and densely populated country like India would rely heavily on distress financing mechanisms to access health care, particularly inpatient care.

From a policy perspective, it is crucial to note that a substantial proportion of household indebtedness in India can be attributed to a general preference for private-sector hospitals (Dilip & Duggal, 2002). Our findings further indicate that indebtedness associated with financing hospitalization care was significantly higher for private healthcare facilities. The data also reveal that the average OOP expenditure in private hospitals is considerably higher than that incurred at public hospitals. These findings are in line with existing literature on in-patient care in India showing that public facilities and publicly funded insurance programmes such as the Rashtriya Swasthya Bima Yojana (RSBY) and PMJAY have demonstrated protective effects against distress modes of financing (Thomas et al., 2024). In light of these observations, it is apparent that the situation of healthcare financing could worsen if the public health system is unable to address the increasing burden of

NCDs in India (Reddy et al., 2005). Such cautionary evidence has emerged from our results, as households seeking care for cancer or CVD are much more likely to incur substantial medical debt (Mahal et al., 2013; Rahman et al., 2013). In this context, although a few social insurance policies (such as Ayushman Bharat, RSBY, and Rajiv Arogyasri Community Insurance Scheme) focus on providing financial protection in India, there remains a lack of comprehensive data on their effectiveness in reducing distressed financing. Nevertheless, given such initiatives, there is a pressing need for investments to expand the public provision of tertiary healthcare, with an exclusive focus on essential diagnostics, drugs and non-medical costs.

The present study, however, is not without limitations, which warrant acknowledgement. First, the non-medical costs associated with treatment-seeking, which can be substantially high, were not captured in the analysis. Additionally, the analysis presented here does not include OOP expenditures related to institutional deliveries. Moreover, several critical but unresolved policy questions remain unaddressed. For instance, there is a lack of information regarding the interest rates and costs associated with borrowing for healthcare expenses. Similarly, the implications of borrowings on basic investments in food and education, as well as the various trade-offs between investing in other competing alternatives, are not well understood. Consequently, deeper insights into coping mechanisms and determinants of successful coping are essential for designing social protection policies, to improve fairness in health care financing.

In conclusion, the heavy reliance on distressed modes of financing, such as borrowings, contributions, and sale of assets, indirectly implies that patients from only affluent sections are reported to afford to access tertiary care, whereas economically vulnerable households might have to compromise with both quantity and quality of care. Furthermore, the findings from this study raise concerns that the situation may worsen with the rising burden of NCDs and the increasing share of aging population.

References

- Bonu, S., Bhushan, I., Rani, M., & Anderson, I. (2009). Incidence and correlates of 'catastrophic' maternal health care expenditure in India. *Health policy and planning*, 24(6), 445-456.
- Bonu, S., Rani, M., Peters, D. H., Jha, P., & Nguyen, S. N. (2005). Does use of tobacco or alcohol contribute to impoverishment from hospitalization costs in India?. *Health Policy and Planning*, 20(1), 41-49.
- Damme, W. V., Leemput, L. V., Por, I., Hardeman, W., & Meessen, B. (2004). Out-of-pocket health expenditure and debt in poor households: evidence from Cambodia. *Tropical Medicine & International Health*, 9(2), 273-280.

Dilip, T. R., & Duggal, R. (2002). Incidence of non-fatal health outcomes and debt in urban India (Draft paper presented at the Urban Research Symposium, December 9–11, 2002). Washington, DC: World Bank.

Erreygers, G. (2009). Correcting the concentration index. *Journal of health economics*, 28(2), 504–515.

Joe, W. (2015). Distressed financing of household out-of-pocket health care payments in India: incidence and correlates. *Health policy and planning*, 30(6), 728–741.

Kruk, M. E., Goldmann, E., & Galea, S. (2009). Borrowing and selling to pay for health care in low-and middle-income countries. *Health Affairs*, 28(4), 1056–1066.

Leckie, G., & Charlton, C. (2013). Runmlwin: a program to run the MLwiN multilevel modeling software from within Stata. *Journal of statistical software*, 52, 1–40.

Leive, A., & Xu, K. (2008). Coping with out-of-pocket health payments: empirical evidence from 15 African countries. *Bulletin of the World Health Organization*, 86(11), 849–856C.

Mahal, A., Karan, A., Fan, V. Y., & Engelgau, M. (2013). The economic burden of cancers on Indian households. *PloS one*, 8(8), e71853.

Ministry of Health & Family Welfare & Government of India. (2017). National Health Policy 2017. <https://nhsrcindia.org/sites/default/files/2021-07/1%20National%20Health%20Policy%202017%20%28English%29%20.pdf>

Narayan D., Patel R., Schafft K., Rademacher A., & Koch-Schulte S. 2000a. *Voices of the Poor: Can Anyone Hear Us?* New York: Oxford University Press.

Narayan, D., Chambers, R., Shah, M. K., & Petesch, P. (2000b). *Voices of the poor: Crying out for change*. World Bank Publications–Books.

NSSO. “Key indicators of social consumption in India – Health, NSS 75th Round 2018”, Ministry of Statistics and Family Planning, Government of India: 2018.

Rahman, M. M., Gilmour, S., Saito, E., Sultana, P., & Shibuya, K. (2013). Health-related financial catastrophe, inequality and chronic illness in Bangladesh. *PloS one*, 8(2), e56873.

Reddy, K. S., Shah, B., Varghese, C., & Ramadoss, A. (2005). Responding to the threat of chronic diseases in India. *The Lancet*, 366(9498), 1744–1749.

Sauerborn, R., Adams, A., & Hien, M. (1996). Household strategies to cope with the economic costs of illness. *Social science & medicine*, 43(3), 291–301.

Sen, G., & Iyer, A. (2012). Who gains, who loses and how: leveraging gender and class intersections to secure health entitlements. *Social science & medicine*, 74(11), 1802–1811.

StataCorp. (2013). Stata statistical software: Release 10. College Station, TX: StataCorp LP.

Thomas, A. R., Muhammad, T., Sahu, S. K., & Dash, U. (2024). Examining the factors contributing to a reduction in hardship financing among inpatient households in India. *Scientific Reports*, 14(1), 7164.

Van Doorslaer, E., O'Donnell, O., Rannan-Eliya, R. P., Somanathan, A., Adhikari, S. R., Garg, C. C., ... & Zhao, Y. (2007). Catastrophic payments for health care in Asia. *Health economics*, 16(11), 1159–1184.

Wagstaff, A., Van Doorslaer, E., & Paci, P. (1991). On the measurement of horizontal inequity in the delivery of health care. *Journal of health economics*, 10(2), 169–205.



Dr. Mayanka Ambade (1991–2025)

In Memory

Sunil Rajpal[#]

Dr. Mayanka Ambade, a rising figure in the fields of health policy, population sciences, and social epidemiology, sadly passed away on 26 January 2025. As an Assistant Professor at the Indian Institute of Technology (IIT) Mandi and an esteemed member of the IHEPA General Council, Dr. Ambade was deeply committed to improving the lives of India's most vulnerable populations through rigorous research. Her academic journey, spanning from the International Institute for Population Sciences to a fellowship at Harvard University, was defined by her insightful analysis of aging, healthcare access, and gender inequality. She leaves behind a legacy of scholarly excellence and is remembered by the academic community as an inspiring researcher, a dedicated mentor, and a remarkably kind human being.

[#] Corresponding Author: Sunil Rajpal, FLAME University, Pune, India
Email: sunil.rajpall@flame.edu.in

The IHEPA community is deeply saddened by the untimely loss of Dr. Mayanka Ambade, Assistant Professor in the Department of Humanities and Social Sciences at the Indian Institute of Technology (IIT) Mandi, who passed away on 26 January 2025. Her sudden departure has left a profound void in the fields of health policy, population sciences, and social epidemiology. Dr. Ambade was an esteemed member of the General Council of the IHEPA and contributed actively to the initiatives with utmost enthusiasm. Despite her young age, Dr. Ambade built a distinguished and impactful body of work, marked by relevant questions, intellectual rigor, remarkable clarity of purpose, and a deep commitment to improving the lives of India's most vulnerable populations.

Born on 9th May 1991, Dr. Ambade graduated in Economics from the University of Mumbai, after which she pursued her Master's and subsequently her PhD in Population Studies at the International Institute for Population Sciences (IIPS), Mumbai. Her doctoral research demonstrated a strong grasp of interdisciplinary methods, blending economics, demography, and public health to produce insights that would characterize much of her later work. Following her PhD, she contributed to several of India's important national health initiatives, working with key datasets such as the National Family Health Survey (NFHS) and the Global Adult Tobacco Survey (GATS).

Dr. Ambade realized the deep significance of socioeconomic gradients in health outcomes early in her career. Her scholarly curiosity took her beyond India, leading to international collaborations across Europe, including work with the University of Lausanne and the Erasmus Mundus Programme. Her postdoctoral journey eventually brought her to the Lakshmi Mittal and Family South Asia Institute at Harvard University, where she served as the inaugural Mittal Institute India Fellow. During this period, she deepened her research on aging and healthcare utilization, examining how socioeconomic disparities shape patterns of morbidity among older adults.

In May 2023, Dr. Ambade joined IIT Mandi as a faculty member, where she quickly established herself as a valued colleague. Her courses on public health and research methodology were appreciated for their clarity and ability to inspire critical thinking. Dr. Ambade particularly enjoyed interacting with students; she was an inspiring teacher and mentor who guided young researchers with patience and enthusiasm. Many students fondly remember her as approachable and deeply invested in their academic growth.

Dr. Ambade's scholarly output reflected commendable depth. Among her notable works was her article in *Economic and Political Weekly*, titled "Economic Growth and Women's Empowerment: A Repeated Cross-sectional Study from India,"¹ which exemplified her ability to tackle complex demographic questions with methodological robustness. Alongside her research, she contributed actively

to the broader professional community, serving as a member of the Early Career Perspectives Panel of the IUSSP and organizing workshops to support early-career scholars.

Beyond academia, Dr. Ambade was passionate about reading, traveling, and experimenting with global cuisines, personal interests that enriched her life and gave her a groundedness that many fondly remember. She possessed an ability to connect with people through genuine empathy and intellectual generosity. Dr. Ambade leaves behind a legacy defined not only by scholarly excellence but also by kindness and integrity. She will be remembered as an inspiring researcher and a remarkable human being whose work and spirit touched many.

References

Ambade, M., & Chattopadhyay, A. (2024). Economic Growth and Women's Empowerment. *Economic & Political Weekly*, 59(36), 73.

Health, Nutrition, and Wellbeing: Evidence, Evaluation, and Way Forward

13th Annual Conference, 12–13 December 2025

Indian Health Economics and Policy Association

Organized by

School of Humanities and Social Sciences, Indian Institute of Technology, Mandi

List of papers selected for Oral Presentation

Technical Session 1A

Md Mohsin, Aligarh Muslim University

Health and Hunger: The Role of Physical, Mental, and Functional Health in Shaping Severe Food Insecurity Among Older Adults in India—Evidence from LASI Wave 1

Kshirabdhii Tanaya Patra, National Institute of Technology Rourkela

Does Social Capital Matter? Insights into Healthy Ageing in India

Aditi, Population Council Consulting P Ltd

Frailty among Older Adults in India: Economic Burden and Insights from LASI

Kripa Nandan, Indian Institute of Technology Roorkee

A Cross-Sectional Study of Healthcare Responsiveness for Migrant and Non-Migrant Middle-Aged and Older Adults in India: Analysis using Longitudinal Ageing Study

Amrendra Kumar Kushwaha, Indian Institute of Technology Mandi

Cooking Fuel, Gendered Exposure, and Lung Function Among Older Adults in India

Technical Session 1B

Raza Mohammad, International Institute for Population Sciences, Mumbai

Age at Childbearing Initiation: A Double-Edged Sword for Child Anthropometric Deficits in South and Southeast Asia

Arpita Shah, Mumbai School of Economics and Public Policy

A multi-scalar spatial analysis of child undernutrition in Gujarat: Districts, Parliamentary Constituencies (PCs) and Villages

Dr Ramesh Athe, Indian Institute of Information Technology Dharwad

Food Fortification to Combat Hidden Hunger in Children: A Systematic Review and Meta-Analysis of Randomized Controlled Trials

Ritika Kapoor, Foundation of Healthcare Technologies Society

Enhancing Nutritional Knowledge and Dietary Practices among Middle School Children: Evidence from Educational Interventions in Delhi and Chandigarh, India

Dr Anjali Dash, Panchayat College, Bargarh

Nutritional variation among children in India: Special reference to Gujarat and Odisha

Technical Session 1C

Abhishek Anand, International Institute for Population Sciences, Mumbai Financial Toxicity Among Gastric and Pancreatic Cancer Survivors in India and Its Impact on Quality of Life

Pubali Das, Department of Economics, Tripura University

In-State vs. Out of State Hospitalization and Catastrophic Health Expenditure Burden for the Patients of Tripura: Evidences from NSS 75th Round Health Data

Dibyendu Biswas, Jindal School of Government and Public Policy, O.P. Jindal Global University, India

Provider Choices and Out-of-Pocket Health Expenditure (OOPE) for Outpatient Care: Evidence From a Multistage Sample Survey in West Bengal, 2024

Ashkar K, Gulati Institute of Finance and Taxation

Does Population Ageing Drive Public Health Expenditure in Indian States? A Dynamic Panel Analysis

Dr Prerana Patil, Indian Council of Medical Research–NIRRH

Out-of-Pocket Expenditure Experienced by Couples Seeking In Vitro Fertilization (IVF) Services at Tertiary Care Facilities in India

Technical Session 2A

Gayathri B, Indian Institute of Technology Mandi

Assessing Cardiovascular Disease Vulnerability in India: A District-Level Index-Based Analysis

Shubham Ranjan, JK Lakshmipat University

Beyond Female Breast Cancer: Gender-Specific Profiles and Outcomes in India

Shruti Hiremath, KLE Academy of Higher Education & Research

Assessing the Temporal Burden of Diabetes Mellitus Attributable to Dietary Risk Factors in India: Insights from Global Burden of Diseases Study: 1990–2021

Priyadarshini Rathore, Manipal University Jaipur

Epidemiology of Cardiovascular Diseases in India: Determinants and Developmental Implications

Shreya Thakur, Quality Council India

Intergenerational Persistence of Smoking Behavior: Evidence From NFHS- 5, 2019-21

Technical Session 2B

Diksha Rani, Indian Institute of Population Sciences, Mumbai

An Analysis of the Age Distribution of Stunting in Countries with Accelerated Reduction Rates, Using Demographic and Health Survey Data

Pramit Chhetri, Sri Sathya Sai Institute of Higher Learning

Water, Sanitation, and Hygiene (WASH) and Malnutrition Among Under-Five Children in India

Dr Archana Singh, RML Awadh University, Ayodhya, Uttar Pradesh

Child Stunting, Caste Discrimination, and Program Effectiveness: Evaluating Heterogeneous Treatment Effects of Integrated Child Development Services in India.

Dr Rudra Narayan Mishra, Gujarat Institute of Development Research, Ahmedabad

Integrated Community-Facility and Policy Frameworks to Combat Malnutrition and Accelerate Development in Aspirational Districts of India

Meenu Bhaskar, Indian Institute of Technology Bombay

The Impact of PMUY on Household Nutrition: Evidence from India

Technical Session 2C

Vishnu Priya V V, Department of Humanities and Social Sciences, Indian Institute of Space Science and Technology

Gendered Access to Information and Public Health Insurance Enrollment Among Poor Households in India

Dr Godwin S K, Government College for Women, Thiruvananthapuram

Moving from Health Insurance to Health Assurance: Kerala's Experience of Contingent Health Security

Prateek Khanna, Amity University

The Road to Universal Health Coverage in India

Sayantani Manna, O P Jindal Global University

Awareness and Utilisation of Health Financing Models: A Study of Insurance Schemes in West Bengal

Shambhavi Mani, Vivekananda College, University of Delhi

Beyond the Frontlines: ASHA workers, Maternal Health and the Social Determinants of Reproductive Wellbeing in Madhya Pradesh.

Technical Session 3A

Ambarish Majumder, ICAFI University, Tripura

Voice-Based Health Technologies for Low-Literacy Populations: A Systematic Review of Accessibility and Effectiveness

Mandeep Bhardwaj, Lovely Professional University

Climate Change, Agricultural Vulnerability, and Public Health Risks in Indian Agricultural Districts: A Spatial Analysis

Pinaki Das, Vidyasagar University

Impact of the ICDS on the Health Status of Pregnant and Lactating Mothers in India: Evidence from NFHS-5

Basant Kumar Panda, Population Council Consulting Pvt. Ltd.

Unlocking the Potential of young women through skilling: Evidence from an evaluation Study in Rajasthan, India

Veeresh Tadahal, Indian Institute of Information Technology, Trichy

Solarization of Primary Health Centers and Health Outcomes: Community Perspectives Using Three Delay Model in Yadgir Aspirational District, Karnataka, India

Technical Session 3B

Shalem Balla, Indian Institute of Technology Mandi

Voltage drops (transitions) in economic empowerment to intrinsic empowerment of women in India

Prateek Singh, Indian Institute of Technology Mandi

Family Transition in Three Decades and Its Effect on Young Married Women's Health in India

Kareena Kaushik, Banasthali Vidyapith

Women's experiences with infertility and the role of assisted reproductive technologies: A qualitative study from Haryana

Juliet F Lalzarzoliani, The ICAFI University, Mizoram

Work-life balance, mental well-being, and family support among working mothers in Aizawl city: A cross-sectional study

Dr Paramasivam P., Takshashila University

Determinants of mental health and wellbeing: evidence from women IT employees in Chennai, Tamil Nadu

Technical Session 3C

Abishek Paul, Institute of Development Studies Kolkata

Does 'Where They Live' Matter? Exploring the Link Between Living Arrangements and Quality of Life Among the Elderly in India

Sasmita Behera, MIT World Peace University, Pune

Does Energy Poverty Influence Healthcare Expenditure? Evidence from Longitudinal Ageing Study in India

Isha Tanwar, Manipal University Jaipur

Impact of Multimorbidity on Labor Force Participation in India: Evidence from LASI

Ali Abbas Rizvi, International Institute for Population Sciences, Mumbai

Family and Elderly Care: Addressing Unmet Needs in Activities of Daily Living Among Older Populations in India

Sabhya Yadav, Birla Institute of Technology and Sciences, Pilani, K K Birla Goa Campus

Cyclicality of Public Health Expenditure Among Indian States: Fiscal and Political Determinants

Technical Session 4A

Tejal Rajaram Varekar, ICMR–National Institute for Research in Reproductive and Child Health

A Systematic Review and Meta–Analysis of Preconception Care Interventions and their Effects on Maternal and Child Health in South–East Asia: Economic and Feasibility Implications for India

Illias K Sheikh, Development Management Institute

Spousal Violence and Risky Birth Spacing Among Indian Women: Insights from NFHS–5

Sachin Saini, Indian Institute of Technology Mandi

Autonomy Under Scrutiny: Women, Abortion Laws and Reproductive Decision Making

Anisha Ojha, IISER, Pune

Pre–Marital Sexual Behaviour Among Women in India: Evidence from NFHS Series

Siddika Banu, Duliajan College

Status of Maternal Health Across the States of India: A Multidimensional Approach

Technical Session 4B

Amal Tomy, Sikkim Manipal Institute of Technology

Digital and Gamified Assessments as Innovations in Adolescent Health and Wellbeing: A Meta–Analytical Review

Debashree Paul, Indian Institute of Technology Kharagpur

Demystifying Adolescent Malnutrition in West Bengal with Ground Realities: Issues and Implications for Policies and Institutions

Kajal Kumari, Central University of Gujarat, Vadodara, Gujarat, India

Navigating Menstrual Equity: A Critical Enquiry into Social Innovation Practices in Kerala, India

Vysagh R K, University of Calicut

Descriptive Profile of eHealth Initiatives in Kerala: Objectives, Implementation and Challenges

Jatinder Singh, Indian Institute of Technology Roorkee

Does Internet Use Shape Lifestyle: Benefits for Healthy Vs Unhealthy Consumption

Technical Session 4C

Dr Sunil Khosla, VIT–AP University

Impact of Financial Inclusion on Multidimensional Health Poverty Among Rural Households in Eastern India

Welisou Mero, Nagaland University

Assessing Public Healthcare Service Quality in Nagaland: Insights from a Case Study in Phek District

Aakriti Aggarwal, Indian Institute of Technology Kanpur

Can Neighbourhood Clinics Deliver? Causal Evidence from Delhi's Mohalla Clinics

Dr Shailendra Kumar, Department of Community Medicine & School of Public Health, PGIMER, Chandigarh

Assessing State–Level Efficiency Using District Hospital Performance Indicators: A Data Envelopment Analysis Approach

Blessy Sarah Mathew, Lovely Professional University, Punjab

Beyond Coverage: Why Ayushman Bharat–Pradhan Mantri Jan Arogya Yojana Struggles to Deliver Healthcare in Jammu and Kashmir

IHEPA**GOVERNING COUNCIL****President**

Manisha Karne

Vice President

Md Zakaria Siddiqui

Secretary

William Joe

Treasurer

Rudra Narayan Mishra

Joint Secretary

Salim Shah

Reshmi Sengupta

Member

Rinshu Dwivedi

Sunil Rajpal

Vinod B Annigeri

About IHEPA

The Indian Health Economics and Policy Association (IHEPA) is a registered society under the Society's Registration Act 1960. The secretariat is located at the International Institute for Population Sciences, Mumbai. IHEPA welcomes all those interested in issues relating to health economics and policy to be a part of this association. IHEPA welcomes young scholars and researchers, grassroots practitioners, the private sector and community - based organizations to participate actively so that all views and experiences can be heard, debated, understood and absorbed.

Aims and Scope

Health Empirics welcomes original research contributions across a broad spectrum of topics related to health economics, healthcare delivery, and health policy assessment. The journal serves as a critical forum for academic researchers, policy analysts, and healthcare leaders, aiming to bridge the gap between rigorous empirical evidence and actionable health policy.

We are committed to publishing original research that demonstrates analytical depth, addressing complex health challenges through robust methodological frameworks.

Key Areas of Inquiry: The scope of the journal encompasses, but is not limited to:

1. **Applications of Health Econometrics:** Advanced statistical methods, causal inference, and quantitative evaluation.
2. **Assessment of Health Policies:** Impact evaluation, cost-effectiveness analysis, and policy design methodologies.
3. **Determinants of Health:** Social, economic, and environmental factors, including risk factor quantification.
4. **Healthcare Management & Financing:** Insurance systems, provider incentives, efficiency analysis, and financial protection.
5. **Inequalities in Health:** Measurement of disparities, equity in access, and distributional analysis.
6. **Methodological Innovations:** Novel data collection, mixed-methods frameworks, and Big Data applications.

Audience: The journal addresses a diverse audience including academic researchers, health policy analysts, healthcare administrators, and international development practitioners.

Editorial Note

Udaya Shankar Mishra

The Health Factor in Multidimensional Poverty: Trends and Inequalities in India, 2005–2021

Komal Ahluwalia, Anirudhan P Edathil, Saroj Kumar and William Joe

Prevalence and Correlates of Overweight and Obesity in Adults and Older Adults in India; Population-Level Estimates Based on Nationally Representative Surveys (2015–21)

Debayanti Bhowmick, Abhishek Kumar and Ajay Kumar Verma

Out-of-Pocket Health Expenditure in India: Level and Composition based on National Sample Survey Data, 2004–05, 2011–12 and 2022

Abhishek Kumar

Livelihood and Status of Tobacco Processing Workers: Insights from Selected States in India

Nayanatara S.Nayak, Rudra N. Mishra, Karabi Mujumdar, Tara Nair and N.L.Narasimha Reddy

Productivity and Technical Change in the Indian Pharmaceutical Sector: A Comparison of Foreign and Domestic Firms

Tulika Rohilla and Boppana Nagarjuna

Burden of Distress Financing for Hospitalization in India: Prevalence and Patterns from Household Health Care Consumption Survey, 2017–18

Sunil Rajpal, Sneha Gupta and Shreya Ronanki

In Memory: Dr. Mayanka Ambade (1991–2025)

Sunil Rajpal